Determination of radioactivity levels of soil samples and the excess of lifetime cancer risk in Rize province, Turkey

S. Dizman¹, F.K. Görür^{2*}, R. Keser¹

¹Abant İzzet Baysal University, Faculty of Sciences and Arts, Department of Physics, 14280, Bolu, Turkey ²Recep Tayyip Erdoğan University, Faculty of Sciences and Arts, Department of Physics, 53100, Rize, Turkey

ABSTRACT

▶ Original article

*Corresponding author:

Dr. F. Korkmaz Görür, Fax: +90 374 253 46 42 E-mail:

filizkorkmaz@yahoo.com

Revised: Sept. 2015 Accepted: Nov. 2015

Int. J. Radiat. Res., July 2016; 14(3): 237-244

DOI: 10.18869/acadpub.ijrr.14.3.237

Background: Turkey, especially the northern part of it, was one of the countries which were contaminated by the Chernobyl accident. Rize is a city located in the Northeastern district of Turkey which was heavily influenced by the Chernobyl nuclear accident. Materials and Methods: In this study, the activity concentrations of natural (226Ra, 232Th, 40K) and artificial (137Cs) were measurements in soil samples collected from 132 different points in Rize province of Turkey using gamma spectrometry with a high-purity germanium detector. *Results:* The average activity concentrations of ²²⁶Ra, ²³²Th and ⁴⁰K were found to be 85.75±11.77, 51.08±9.42, and 771.57±37.65 Bq/kg in soil samples, respectively. In order to evaluate the radiological hazard of the natural radioactivity, radium equivalent activity (Ra_{eq}), representative level index (I_{vr}), the external hazard index (Hex), the total absorbed dose rate (D), the annual effective dose equivalent (AEDE) and excess life time cancer risk (ELCR) have been calculated and compared with the internationally approved values. Conclusion: The outdoor air absorbed dose rates (D) due to terrestrial gamma rays for soil have been calculated because of agricultural area and living in the surrounding. It is important to determine background radiation level in order to evaluate the health hazards. Annual effective gamma doses and the lifetime risks of cancer were higher than the world's average. Moreover compared to the World's average, the lifetime risk of cancer doubled for most of the localities.

Keywords: Radioactivity, soil, lifetime cancer risk, gamma dose.

INTRODUCTION

Human beings are exposed to background radiation that stems both from natural and manmade sources. Natural background radiation, which is equivalent to 2.4 mSv per person, makes up approximately 80% of the total radiation dose a person is exposed in a year (1). Soil radionuclide activity concentration is one of determinants of the main natural radiation. Volcanic geographic background structures as well as rocks that are rich in phosphate, granite and salt contain natural radionuclides like ²³⁸U, ²³²Th and ⁴⁰K. When disintegrated through

processes, radionuclides are carried to the soil by rain and flows ⁽²⁾.

The natural radioactivity in soil comes from ²²⁶Ra, ²³²Th and from natural ⁴⁰K. Some other terrestrial radionuclides, including those of the ²³⁵U series, ⁸⁷Rb, ¹³⁸La, ¹⁴⁷Sm and ¹⁷⁶Lu exist in nature but at such low levels that their contributions to the dose in the humans are small. Artificial radionuclides can also be present such as ¹³⁷Cs, resulting from fallout from weapons testing. ¹³⁷Cs is a fission product which is formed through nuclear tests and accidents. The deposition of ¹³⁷Cs in soil is important since its half-life is 30.2 years and it has a gamma emission of 661 keV ⁽³⁾. The radiological

Dizman et al. / Radioactivity levels of soil and lifetime cancer risk

implication of these radionuclides is due to the gamma-ray exposure of the body and irradiation of lung tissue from inhalation of radon and its daughters. Therefore, the assessment of gamma radiation dose from natural sources is of particular importance as natural radiation is the largest contributor to the external dose of the world population (4). Natural environmental radioactivity and the associated external exposure due to gamma radiation depends primarily on the geological and geographical conditions, and appear at different levels in the soils of each region in the world (5).

Turkey, especially the northern part of it, was one of the countries which were contaminated by the Chernobyl accident (6). Rize is a city located in the Northeastern district of Turkey which was heavily influenced by the Chernobyl nuclear accident. The radioactive plume from the accident reached Turkey by 5 May 1986, substantially contaminating various regions and ecosystems of the country. During the emergency, Cekmece Nuclear Research and Training Center (CANEM) performed an analysis of various substances. It their report, it has been noted that the surface soil 137Cs activity concentration of the eastern part of the Black Sea mountains was around 4000-4500 Bq/kg at the 0.5cm soil in the year 1988 (7).

It is critical to evaluate soil radioactivity in order to understand background radiation

concentrations. Measuring terrestrial gamma dose rates is also essential since gamma radiation provides information concerning excess lifetime cancer risks. Yet in Turkey, there are only a limited number of studies which evaluate soil radioactivity and terrestrial gamma dose rates (8-25).

The aim of this study is to determine natural (226Ra, ²³²Th, ⁴⁰K) and artificial radioactivity levels in soil collected from different points in Rize province of Turkey. Also, the average radium equivalent activity (Ra_{eq}), representative level index (I_{yr}) , the external hazard index (Hex), the total absorbed dose rate (D), the annual effective dose equivalent (AEDE) and excess life time cancer risk (ELCR) which will be defined later have been calculated and compared with the results in literature. The results of this study will provide background data on the natural and artificial radioactive isotopes and environmental pollution.

MATERIALS AND METHODS

Study area

Rize is a province of north-east Turkey, on the eastern Black Sea coast (figure 1). Rize stands between the latitudes of 40°-20' and 41°-20' N and the longitudes of 40°-22' and 41°-28' E. Rize has a catchment area of 3920 km². It is on the




Figure 1. Location of sampling sites indicating the Rize Province, Turkey.

north side of the range of mountains that run along the Black Sea coast. Overlooking the sea this is the wettest corner of Turkey and Rize is the country's largest producer of tea. Summers are cool (July average 22°C), winters are warm (January average 7°C) and it is wet all year round. It has twelve district areas. The population of Rize is 361353 (26).

Sampling and sample preparation

Soil samples were collected from 12 locations of the study area during the year 2010. In all a total of 132 samples were analyzed. After clearing the ground of stones, pebbles, vegetation and roots, 1-2 kg of material from the first 10 cm of topsoil was placed in labeled polythene bags and then transferred to the laboratory. The samples were dried at 60 °C for 48 h, grained, passed through 2 mm sieves. The dried samples then were homogenized and weighed and transferred into uncontaminated empty cylindrical plastic containers of uniform size. The samples were weighed and stored for a minimum period of one month to allow daughter products to come into radioactive secular equilibrium with their parents ²²⁶Ra and ²³²Th and then were counted for 50.000-100.000 s the concentration of the depending on radionuclides.

Experimental method for γ- spectroscopy

The radiation levels of samples were analyzed using gamma spectrometry, which was equipped with a 55% efficiency high purity germanium (HPGe) detector and a multi-channel analyzer. The gamma spectra were analyzed by using the ORTEC Maestro 32 data acquisition and analysis system. The detector had coaxial closed–facing geometry with the following specifications: resolution full width half maximum (FWHM) at 122 keV ⁵⁷Co was 1.00 keV and at 1.33MeV ⁶⁰Co was 1.90 keV. The detector was shielded by a cylindrical lead shield, which had average thickness of 10 cm in order to achieve a background level as low as possible.

Efficiency of the detector was determined with a ¹⁵²Eu source (Amersham Company, UK) of known activity. ¹⁵²Eu source have been widely

used for calibration and efficiency determination due to their large range of energies (122, 244, 344, 411, 443, 779, 964, 1112 and 1408 keV) with emission probabilities of 3-29 % (27, 28). An ideal measuring geometry of cylindrical source (homogeneously distributed activity with constant volume and distance) was placed coaxially with the detector for the efficiency determination and the same procedure applied for the sample measurements.

Soil samples were placed symmetrically on top of the detector and measured for a period of 100.000 s. The net area under the corresponding peaks in the energy spectrum was computed by subtracting counts due to Compton scattering of higher peaks and other background sources from the total area of the peaks. From the net area of a certain peak, the activity concentrations in the samples were obtained using equation 1:

$$C(Bq/kg) = \frac{C_n}{\epsilon P_{\gamma} M_s}$$
 (1)

where C is the activity concentration of the radionuclide in the sample given in Bq/kg, C_n is the count rate under the corresponding peak, ϵ is the detector efficiency at the specific γ -ray energy, P_{γ} is the absolute transition probability of the specific γ -ray, and M_S is the mass of the sample (kg).

For the measurement of 226 Ra activity concentration, the -ray energies of 295.21 and 351.92 keV of 214 Pb, 609.31 keV of 214 Bi were used. The activity concentration of 232 Th was determined at the γ -ray energies 911.07 keV and 969.11 keV of 228 Ac, 40 K and 137 Cs were measured directly from the 1460.8 keV and 661.66 keV peak energies, respectively $^{(29,30)}$. 40 K activity determined from the 1460.7 keV emission gamma-lines and 137 Cs activity determined from the 661.1 keV emission gamma-lines.

Calculation of the radiological effects

The activity concentrations of ²²⁶Ra, ²³²Th, ⁴⁰K and ¹³⁷Cs measured in each of the soil samples. Radium equivalent activity is a widely used hazard index and it is calculated through the relation given by Beretka and Mathew ⁽³¹⁾. It is assumed that 370 Bq/kg of ²²⁶Ra, 259 Bq/kg of

Int. J. Radiat. Res., Vol. 14 No. 3, July 2016

 232 Th and 4810 Bq/kg of 40 K produce the same gamma-ray dose rate

$$Ra_{eq} = A_{Ra} + 1.43A_{Th} + 0.077A_{K}$$
 (2)

where A_{Ra} , A_{Th} and A_{K} are the activity concentration of 226 Ra, 232 Th and 40 K in Bq/kg, respectively.

Another radiation hazard index called the representative level index, $I_{\gamma T}$, is defined from the following formula (32, 33):

$$I_{\gamma r} = \frac{1}{150Bq/kg} A_{Ra} + \frac{1}{100Bq/kg} A_{Th} + \frac{1}{1500Bq/kg} A_{K}$$
 (3)

where A_{Ra} , A_{Th} , and A_{K} have the same meaning as in equation 2.

The external hazard index, H_{ex} was calculated for the investigated samples using the model proposed by Krieger $^{(34)}$ assuming thick walls without windows and doors, where the external hazard index is given by

$$H_{ex} = A_{Ra}/370 + A_{Th}/259 + A_{K}/4810 \le 1$$
 (4)

where A_{Ra} , A_{Th} and A_{K} are the activity concentration of 226 Ra, 232 Th and 40 K in Bq/kg, respectively. The calculated average external hazard index was found to be less than unity.

The absorbed dose rate, D (nGy/h) in air at 1 m above ground level due to the presence of 226 Ra, 232 Th, 40 K and 137 Cs in the soil samples at each site was calculated using the following equation 5 $^{(5)}$,

$$D=aC_{Ra}+bC_{Th}+cC_{K}+dC_{Cs}$$
 (5)

where a is the dose rate per unit ²²⁶Ra activity concentration (4.27x10⁻¹⁰ Gy/h/Bq/kg), C_{Ra} is the concentration of ²²⁶Ra in the sample (Bq/kg), b is the dose rate per unit ²³²Th activity concentration (6.62x10-10 Gy/h/ Bq/kg), C_{Th} is the concentration of ²³²Th in the sample (Bq/ kg), c is the dose rate per unit 40K activity concentration (0.43x10⁻¹⁰ Gy/h/ Bq/kg), C_K is the concentration of 40K in the sample (Bq/kg), d is the dose rate per unit ¹³⁷Cs activity concentration (0.30 x10⁻¹⁰ Gy/h/ Bq/kg) and C_{Cs} is the concentration of ¹³⁷Cs in the sample (Bq/ kg). The absorbed dose rate (nGy/h) in air at 1 m above the ground determined at each farm does not directly give the radiological hazard to which an individual is exposed.

The annual effective dose equivalent (AEDE) was calculated by using equation 6:

AEDE
$$(\mu Sv/y) = D \times DCF \times OF \times T$$
 (6)

where D is absorbed dose rate in air (nGy/ h), DCF is dose conversion factor (0.7 Sv/Gy), OF is outdoor occupancy factor (0.2), T is the time $(8760 \text{ h/y})^{(5)}$.

Excess life time cancer risk (ELCR) was calculated by using equation 7:

$$ELCR = AEDE \times DL \times RF$$
 (7)

where DL is duration of life (70 years) and RF is risk factor (Sv⁻¹). For stochastic effects, ICRP 90 uses values of 0.05 for the public ⁽²⁹⁾.

RESULTS AND DISCUSSION

The results of activity concentrations in the soil samples from twelve different sites are gives in table 1 for the natural radionuclides of 226 Ra, 232 Th and 40 K and the artificial radionuclide of 137 Cs. Radium equivalent activity (Ra_{eq}), representative level index, external hazard index, absorbed dose rates, annual effective doses and the excess lifetime risks of cancer in soil samples are given in table 2.

The concentrations found in the present study ranged from 48.54±6.30 (İyidere district) to 163.14±21.29 (Kalkandere district) Bq/kg for ²²⁶Ra, from 19.58±4.01 (Hemsin district) to 125.53±22.74 (Güneysu district) Bq/kg for ²³²Th, from 302.40±15.48 (Güneysu district) to 1159.51±61.06 (Hemşin district) Bq/kg for ⁴⁰K. The average activity concentrations of ²²⁶Ra, ²³²Th and ⁴⁰K were found to be 85.75±11.77, 51.08±9.42, and 771.57±37.65 Bq/kg in soil samples, respectively. The activity of ⁴⁰K is seen to be higher than 226Ra and 232Th in all the selected soil samples. The world's mean values of ²²⁶Ra, ²³²Th and ⁴⁰K activity concentrations are 32, 45 and 420 Bq/kg, respectively (5). The mean values of ²²⁶Ra, ²³²Th and ⁴⁰K are higher than the world's average values.

¹³⁷Cs does not exist in soil naturally. It is a product of fallout radioactivity. The ¹³⁷Cs might have been deposited in soil of study area, presumably as a result of the nuclear power

plant accident at Chernobyl on 26 April 1986. Moreover, measured ¹³⁷Cs activity concentrations can be attributed to the atmospheric nuclear weapon tests conducted by several countries. ¹³⁷Cs activities in soil samples varied from 75.80±6.30 (İyidere district) to 481.81±30.07 (Güneysu district) Bq/kg and average ¹³⁷Cs activity was found to be 236.38±13.49 Bq/kg.

As shown in table 3, the radioactivity concentrations in soil samples were comparable

to other studies in various regions (8-25,35-40). Karadeniz *et al.*, Tabat *et al.*, Kiliç *et al.*, Kam and Bozkurt, Celik *et al.* determined slightly lower activity concentrations of ²²⁶Ra and ²³²Th compared to this study (8,12,17,19,25). Merdanoğlu and Altınsoy *et al.*, Orgun *et al.* and Abbaspour *et al.* determined a higher activity concentrations of ²²⁶Ra and ²³²Th compared to this study (13,20,35). ⁴⁰K activity concentrations of Rize were also higher compared to studies conducted at other parts of Turkey (8,12,14-17,19,21,22).

Table 1. Radioactivity concentrations of ²²⁶Ra, ²³²Th, ⁴⁰K and ¹³⁷Cs in soil samples.

| | Number of sampling | ²²⁶ Ra (Bq/kg) | ²²⁶ Ra (Bq/kg) | | ¹³⁷ Cs (Bq/kg) | |
|----------------------|--------------------|---------------------------|---------------------------|---------------|---------------------------|--|
| Rize Centrum | 12 | 66.50±11.54 | 48.97±8.29 | 842.38±45.64 | 240.32±6.99 | |
| Derepazarı district | 12 | 79.40±12.53 | 44.49±8.40 | 876.98±45.70 | 118.26±5.11 | |
| Ardeşen district | 12 | 58.19±11.37 | 24.29±6.10 | 654.25±38.54 | 244.15±6.82 | |
| Pazar district | 12 | 65.57±11.11 | 28.29±5.72 | 645.94±35.27 | 374.18±7.78 | |
| Fındıklı district | 12 | 50.99±11.44 | 26.59±6.94 | 627.95±41.15 | 322.84±8.65 | |
| Ikizdere district | 10 | 92.28±9.65 | 47.25±5.43 | 871.72±34.11 | 87.68±4.99 | |
| Kalkandere district | 12 | 163.14±21.29 | 89.57±17.32 | 708.62±33.82 | 376.69±18.18 | |
| Çayeli district | 12 | 149.64±22.06 | 68.24±15.54 | 634.78±36.35 | 317.58±37.31 | |
| Çamlıhemşin district | 8 | 67.86±10.82 | 24.71±7.39 | 810.78±28.21 | 82.49±9.73 | |
| Hemşin district | 8 | 101.17±14.46 | 19.58±4.01 | 1159.51±61.06 | 114.78±20.00 | |
| Güneysu district | 10 | 85.67±10.34 | 125.53±22.74 | 302.40±15.48 | 481.81±30.07 | |
| lyidere district | 12 | 48.54±6.30 | 65.40±5.20 | 1123.54±36.57 | 75.80±6.30 | |
| Total district | 132 | 85.75±11.77 | 51.08±9.42 | 771.57±37.65 | 236.38±13.49 | |

Table 2. Radium equivalent, representative level index, external hazard index, absorbed dose rates, annual effective doses and the excess lifetime risks of cancer in soil samples in Rize.

| | Ra _{eq} (Bq/kg) | l _{γr} | H _{ex} | D (nGy/h) | AEDE (mSv/y) | Life time total dose(mSv) | ELCR (×10 ⁻³) |
|----------------------|--------------------------|-----------------|-----------------|--------------|-----------------|------------------------------|---------------------------|
| Rize Centrum | 201.39 | 1.49 | 0.54 | 104.25 | 0.127 | 7.30 | 0.45 |
| Derepazarı district | 210.55 | 1.56 | 0.57 | 104.61 | 0.128 | 7.32 | 0.45 |
| Ardeşen district | 143.30 | 1.07 | 0.39 | 76.38 | 0.093 | 5.35 | 0.33 |
| Pazar district | 155.76 | 1.15 | 0.42 | 85.73 | 0.105 | 6.00 | 0.37 |
| Fındıklı district | 137.37 | 1.02 | 0.37 | 76.06 | 0.093 | 5.32 | 0.33 |
| Ikizdere district | 226.97 | 1.67 | 0.61 | 110.80 | 0.135 | 7.76 | 0.48 |
| Kalkandere district | 345.79 | 2.46 | 0.93 | 170.73 | 0.209 | 11.95 | 0.73 |
| Çayeli district | 296.10 | 2.10 | 0.80 | 145.89 | 0.178 | 10.21 | 0.63 |
| Çamlıhemşin district | 165.63 | 1.24 | 0.45 | 82.67 | 0.101 | 5.79 | 0.35 |
| Hemşin district | 218.45 | 1.64 | 0.59 | 109.46 | 0.134 | 7.66 | 0.47 |
| Güneysu district | 288.46 | 2.03 | 0.78 | 147.14 | 0.180 | 10.30 | 0.63 |
| lyidere district | 228.57 | 1.73 | 0.62 | 114.61 | 0.140 | 8.02 | 0.49 |
| Total district | 218.20 | 1.60 | 0.59 | 110.69 | 0.136 | 7.75 | 0.48 |
| World ⁽⁵⁾ | - | | <1 | 60 | 0.070 | 4.90 | 0.29 |

Dizman et al. / Radioactivity levels of soil and lifetime cancer risk

Table 3. ²²⁶Ra, ²³²Th, ⁴⁰K and ¹³⁷Cs activity concentrations, absorbed dose rates (D), annual effective dose equivalents (AEDE) and excess lifetime risks of cancer (ELCR) in various studies.

| Area | ²²⁶ Ra (Bq/kg) | ²³² Th (Bq/kg) | ⁴⁰ K (Bq/kg) | ¹³⁷ Cs (Bq/kg) | D (nGy/h) | AEDE (μSv/y) | ELCR (x10-3) |
|----------------------------|------------------------------|------------------------------|----------------------------|------------------------------|--------------|-----------------|-----------------|
| Rize (Present study) | 85.75 | 51.08 | 771.57 | 236.38 | 110.69 | 136 | 0.48 |
| Kırklareli ⁽¹⁴⁾ | 37 | 40 | 667 | 8 | 71 | 87 | 0.51 |
| Manisa ⁽¹²⁾ | - | 27 | 340 | - | 54 | 66 | - |
| Istanbul ⁽¹⁶⁾ | - | 37 | 342 | 1.8-81 | 49 | 65 | - |
| Marmara ⁽¹⁷⁾ | 22.5 | 26.6 | 443 | 0.9-154 | 44.73 | 54.86 | - |
| Kestanbol (18) | 130.93 | 192 | 1207 | 0.37-36 | 219 | 269 | - |
| Kastamonu ⁽¹⁹⁾ | 37.4 | 27.17 | 431.43 | 8.02 | 52.76 | 65 | - |
| Çanakkale ⁽²⁰⁾ | - | 204.69 | 1171 | 0-6.57 | 178-448 | - | - |
| Kazdağı ⁽⁸⁾ | 21.7 | 21.1 | 297.5 | 0.1-28 | 44 | 67 | - |
| Sanliurfa ⁽²²⁾ | - | 24.95 | 298.6 | 9.08 | 38.24 | 60.09 | - |
| Fırtına ⁽²³⁾ | 15-116 | 10-105 | 105-1235 | 19-232 | 77.4 | - | - |
| Eastern Black Sea (24) | 12-120 | 40.9 | 622.8 | 169.7 | 77.18 | 93.30 | - |
| Giresun (25) | 33 | 43 | 733 | 318 | - | 92 | - |
| Iran ⁽³⁵⁾ | 1188 | 64.92 | 545.10 | 10.41 | 612.37 | 750 | 0.30 |
| India ⁽³⁶⁾ | 57.34 | 52.83-105.81 | 95.33-160.30 | - | 72.35-108.65 | 72.7-133.2 | - |
| China ⁽³⁷⁾ | - | 71.5 | 672 | - | 124 | 152 | - |
| India ⁽³⁹⁾ | 50.58 | 63.13 | 268.92 | - | 66.89 | 490 | - |
| Nigeria ⁽⁴⁰⁾ | - | - | 1190 | 1 | 52-414 | 200 | - |
| Worldwide ⁽⁵⁾ | 35 | 45 | 420 | - | 60 | 70 | 0.25 |

The values of Ra_{eq} varied from 137.37 to 345.79 Bq/kg and average value of Ra_{eq} was found to be 218.20 Bq/kg. The estimated average values of Ra_{eq} in the present work are lower than the recommended maximum value of 370 Bq/kg ⁽³¹⁾. On comparing the measured mean values from some of the other studies, it is observed that value of this work is higher than the measured values of 166.3 Bq/kg in Firtina Valley ⁽²³⁾. The values of representative level index, $I_{\gamma r}$ varied from 1.02 to 2.46 and average value of $I_{\gamma r}$ was found to be 1.60.

The values of external hazard index, H_{ex} range from 0.37 to 0.93 and average value was found to be 0.59 for the soil samples. The maximum value of H_{ex} must be less than unity. All values estimated of H_{ex} in the present work are lower than unity. The average values of H_{ex} were found to be 0.50 for India, 0.45 for Firtina Valley of Turkey and 0.84 for Xiazhuang Granite Area (China) $^{(36,23,37)}$.

The absorbed dose rates in air for soil samples the average dose rate was 110.69 nGy/h in Rize. The average D value for soil was

calculated as 71, 77.4, 77.18, 44, 178-448 nGy/h in Kırklareli, in Adana, in Çanakkale, in Fırtına Valley, in Eastern Black Sea, respectively (14,20,21,23,24). The population weighted values give an average absorbed dose rate in air outdoors from terrestrial gamma radiation of 60 nGy/h (5). This reveals that the mean absorbed dose rate in air outdoors from Rize areas is almost two times higher than that of the worldwide average value.

As shown in table 2 and 3, the calculated values of annual effective dose for the all soil samples ranged from 93 to 209 μ Sv/y, with a mean of 136 μ Sv/y, which is higher than the world average value of 70 μ Sv/y ⁽⁵⁾. So, the obtained values are higher than the world average value. The average AEDE value was calculated to be 87 μ Sv/y in Kırklareli, 66 μ Sv/y in Manisa, 65 μ Sv/y in the Istanbul (Turkey), 60.09 μ Sv/y in Sanliurfa ^(14-16,22). These average values are generally lower than our result.

As shown in table 2, when life expectancy was taken as 70 y, the average life time total gamma radiation was calculated as 7.75 mSv, which yielded a lifetime cancer risk of 0.48×10^{-3} . The

world's mean value of life time total gamma dose and the excess lifetime cancer risk are 4.90 mSv and 0.29×10⁻³, respectively ⁽⁵⁾. The mean of life time total gamma dose and the excess lifetime cancer risk observed in this study are higher than the world's mean values. The average ELCR value was calculated to be 0.5×10⁻³ in Kırklareli, 0.26×10-3 in western Mazandaran (Iran) (14,35). Yet, due to the unavailability of related mortality and morbidity statistics, the health hazards of the assessed values on the population were not calculated. Therefore, this study was limited to background radiation levels.

CONCLUSION

The obtained data cover a wide area in Rize. The mean concentrations of the radionuclides ²²⁶Ra, ²³²Th, ¹³⁷Cs, and ⁴⁰K in soil samples determined in this study compare suitably with literature values. But the ¹³⁷Cs activity concentrations in some places are higher than the other results. This can be attributed to the Chernobyl nuclear power plant accident and the atmospheric nuclear weapon tests conducted by several countries. From the measured values. the average values of radium equivalent activity (Ra_{eq}), representative level index (I_{yr}), external hazard index (H_{ex}), absorbed dose rate in air (D), annual effective dose equivalent (AEDE) and the excess life time cancer risk (ELCR) were calculated. The outdoor air absorbed dose rates (D) due to terrestrial gamma rays for soil have been calculated because of agricultural area and living in the surrounding. It is important to determine background radiation level in order to evaluate the health hazards. Annual effective gamma doses and the lifetime risks of cancer were higher than the world's average. Moreover compared to the World's average, the lifetime risk of cancer doubled for most of the localities.

Conflict of interest: Declared None

REFERENCES

1. IAEA (International Atomic Energy Agency) (1996)

Technical Report Series 96-00725, Vienna, Austria.

- 2. NCRP (National Council on Radiation Protection and Measurement) (1975) Natural background radiation in the U.S., Soil Radioactivity, No. 45.
- 3. Teodere DF and Edwards JN (1993) Housing, stres and well-being, evidence from Thailand. Soc Sci Med, 36: 1417
- 4. UNSCEAR (United Nations Scientific Committee on the Effects of Atomic Radiation) (1988) Sources and Effects of Ionizing Radiation, United Nations, New York.
- 5. UNSCEAR (2000) Sources and effects of ionizing radiation, Vol. 1. United Nations Scientific Committee on the Effects of Atomic Radiation. Report of the General Assembly with Scientific Annexes. United Nations, New York.
- 6. TAEK (1998) Türkiye'de Çernobil sonrası Radyasyon ve Radyasyon çalışmaları, TAEK report, April 1998.
- 7. Unlu MY, Topcuoglu S, Kucukcezzar R, Varinlioglu A, Gungor N, Bulut AM, Gungor E (1995) Natural effective half-life of ¹³⁷Cs in tea plants. Health Phys, 65: 94–99.
- Karadeniz O, Karakurt H, Akal C (2015) Natural radionuclide activities in forest soil horizons of Mount IDA/ Kazdagi, Turkey. Environ Monit Assess, 187(6): 319.
- Agar O, Boztosun I, Korkmaz ME, Ozman SF (2014) Measurement of Radioactivity Levels and Assessment of Radioactivity Hazards of Soil Samples in Karaman, Turkey. Radiat Prot Dosim, 162: 630-637.
- 10. Yıldız N, Oto B, Turhan Ş, Uğur FA, Gören E (2014) Radionuclide determination and radioactivity evaluation of surface soil samples collected along the Erçek Lake basin in eastern Anatolia, Turkey. J Geochem Explo, 146: 34-39.
- 11. Damla N and Aldemir K (2014) Radon survey and soil gamma doses in primary schools of Batman, Turkey. Isotopes Environ Health, 50: 226-234.
- 12. Tabar E, Kumru MN, Ichedef M, Sac MM (2013) Radioactivity level and the measurement of soil gas radon concentration in Dikili geothermal area, Turkey. Int J Radiat Res, 11(4): 253-261.
- 13. Keser R, Gorur FK, Akcay N, Okumusoglu NT (2011) Radionuclide concentration in tea, cabbage, orange, kiwi and soil and lifetime cancer risk due to gamma radioactivity in Rize, Turkey. J Sci Food Agr, 91: 987-991.
- 14. Taskin H, Karavus M, Ay P, Topuzoglu A, Hidiroglu S, Karahan G (2009) Radionuclide concentrations in soil and lifetime cancer risk due to gamma radioactivity in Kirklareli, Turkey. J Environ Radioactiv, 100: 49-53.
- 15. Erees FS, Akozcan S, Parlak Y, Çam S (2006) Assessment of dose rates around Manisa (Turkey). Radiat Meas, 41: 598-
- 16. Karahan G and Bayulken A (2000) Assessment of gamma dose rates around Istanbul(Turkey). J Environ Radioact, 47: 213-221.
- 17. Kiliç O, Belivermis M, Topçuoglu S, Cotuk Y, Coskun M, Cayir A, Kuçer R (2008) Radioactivity concentrations and dose assessment in surface soil samples from east and south of Marmara Region, Turkey. Radiat Prot Dosim, 128 (3): 324-330.
- 18. Merdanoglu B and Altinsoy N (2006) Radioactivity

Int. J. Radiat. Res., Vol. 14 No. 3, July 2016

243

Dizman et al. / Radioactivity levels of soil and lifetime cancer risk

- concentrations and dose assessment for soil samples from Kestanbol granite area, Turkey. *Radiat Prot Dosim*, **121**: 399–405.
- Kam E and Bozkurt A (2007) Environmental radioactivity measurements in Kastamonu region of northern Turkey. Appl Radiat Isot, 65: 440–444.
- Orgun Y, Altınsoy N, Sahin SY, Gungor Y, Gultekin AH, Karahan G, Karacık Z (2007) Natural and anthropogenic radionuclides in rocks and beach sands from Ezine region (Canakkale), Western Anatolia, Turkey. Appl Radiat Isot, 65: 739–747.
- Degerlier M, Karahan G, Ozger G (2008) Radioactivity concentrations and dose assessment for soil samples around Adana, Turkey. J Environ Radioact, 99: 1018–1025.
- Bozkurt A, Yorulmaz N, Kam E, Karahan G, Osmanlioglu AE (2007) Assessment of environmental radioactivity for Sanliurfa region of southeastern Turkey. *Radiat Meas*, 42: 1387–1391.
- Kurnaz A, Kuçukomeroglu B, Keser R, Okumusoglu NT, Korkmaz F, Karahan G, Cevik U (2007) Determination of radioactivity levels and hazards of soil and sediment samples in Firtina Valley (Rize, Turkey). Appl Radiat Isot, 65: 1281–1289.
- Celik N, Cevik U, Celik A, Koz B (2009) Natural and artificial radioactivity measurements in Eastern Black Sea region of Turkey. J Hazard Mater, 162: 146–153.
- Celik N, Cevik U, Celik A, Kucukomeroglu B (2008) Determination of indoor radon and soil radioactivity levels in Giresun, Turkey. J Environ Radioactiv, 99: 1349–1354.
- ITS Turkish Statistical Institute (2007) News Bulletin 9, T.C.
 Bas bakanlık Turkiye İstatistik Kurumu, Turkey.
- 27. Firestone RB and Shirley VS (1998) Table of Isotopes, 8th Edition, John Wiley, New York.
- Grigorescu EL, Razdolescu AC, Sahagia M, Luca A, Ivan C, Tanase G (2002) Standardization of ¹⁵²Eu. Appl Radiat Isot, 56: 435-439.
- 29. ICRP (International Commission on Radiological Protection) (1990) Recommendations of the International

- Commission on Radiological Protection, Vol. 21, No.1–3, Publication 60.
- IAEA (International Atomic Energy Agency) (1989) Vienna, Austria, Technical Report Series-295.
- Beretka J and Mathew PJ (1985) Natural radioactivity of Australian building materials, industrial wastes and byproducts. Health Phys, 48: 87-95.
- 32. Alam MN, Chowdhury MI, Kamal M, Ghose S, Ismal MN (1999) The ²²⁶Ra ²³²Th and ⁴⁰K activities in beach sand minerals and beach soils of Cox'sBazar, Bangladesh. *J Environ Radioactiv*, **46 (2):** 243–250.
- 33. NEA-OECD (Nuclear Energy Agency) (1979) Exposure to radiation from natural radioactivity in building materials. Report by NEA Group of Experts OECD, Paris.
- 34. Krieger (1981) Radioactivity of construction materials, Betonwerk Fertigteil Techn. 47, 468.
- 35. Abbaspour M, Moattar F, Okhovatian A, Kharrat SM (2010) Relationship of Soil Terrestrial Radionuclide Concentrations and The Excess of Lifetime Cancer Risk in Western Mazandaran Province, Iran. Radiat Prot Dosim, 142 (2-4): 265-272.
- Rani A and Singh S (2005) Natural radioactivity levels in soil samples from some areas of Himachal Pradesh, India using g-ray spectrometry. Atmos Environ, 39: 6306–6314.
- Yang Y, Wu X, Jiang Z, Wang W, Lu J, Lin J, Wang L, Hsia Y (2005) Radioactivity concentrations in soils of the Xiazhuang granite area, China. Appl Radiat Isot, 63: 255–259
- 38. Rafique M (2014) Cesium-137 activity concentrations in soil and brick samples of Mirpur, Azad Kashmir; Pakistan. *Int J Radiat Res,* **12 (1):** 39-46.
- 39. Dhawal SJ, Kulkarni GS, Pawar SH (2013) Terrestrial background radiation studies in South Konkan, Maharashtra, India. *Int J Radiat Res*, **11 (4)**: 263-270.
- Ajayi OS and Ibikunle SB (2013) Radioactivity of surface soils from Oyo state, South Western Nigeria. Int J Radiat Res, 11 (4): 271-278.