Development and statistical assessment of a radiation safety literacy measurement tool

K. Choi and J.K. Cho*

Department of Radiological Science, Jeonju University, Jeonju, Republic of Korea

ABSTRACT

Background: The public's understanding of radiation and protection of radiation harm is not high. Therefore, it is meaningful to know the radiation safety literacy level in terms of health care. This study developed a measurement tool that can help to identify the actual condition of radiation safety literacy and conducted the statistical assessment of the developed tool. Materials and Methods: We developed a radiation safety literacy measurement tool in four steps: preliminary term extraction, content validity, face validity, and response scale composition using expert groups such as radiologist, professor of radiological science, angiography nurse, and professor of Korean language. And we developed a questionnaire and conducted a survey on samples of n = 280 (male: 124, female: 156). For statistical assessment, descriptive analysis, Cronbach's coefficient, and correlation analysis were performed, and receiver operating characteristic (ROC) curve was obtained. Results: As a result of developing measurement tool of four steps, radiation safety literacy measurement tool consisting of a total of 46 items were developed. The result of the survey showed high reliability with the internal consistency reliability coefficient of 0.963. The correlation coefficient of the developed measurement tool with the rapid estimate of adult literacy in medicine was found to be valid with 0.448 (p<0.05). Conclusion: The radiation safety literacy measurement tool developed in this study can be used as a useful tool for the process of patient evaluation for appropriate communication between the healthcare provider and the patient regarding radiation.

Keywords: Radiation safety, health literacy, reliability, validity, receiver operating characteristic curve.

Original article

*Corresponding authors: Jung Keun Cho, Ph.D.,

E-mail: cjk0129@naver.com

Revised: November 2019 Accepted: January 2020

Int. J. Radiat. Res., January 2021;

19(1): 41-48

DOI: 10.29252/ijrr.19.1.41

INTRODUCTION

Every human being wants to live long and healthy. There are many factors that affect health, and recent studies have shown that health information literacy is also one of factors. Health information literacy is defined as the ability to use and understand health-related information (1). In Korea, it is given different names "medical information such as comprehension ability," "medical information literacy," "health information utilization ability," "health literacy," "ability to understand health information," and "health literacy" (2). "Ability to understand health information" includes the ability of individuals to understand explanations from medical staff when using medical care, reading or filling out health questionnaires and consent forms, understanding health education materials, and solving descriptions of basic medication and methods (3). To sum up the above, the health information literacy can be defined as "the ability to acquire basic health information and a series of processes to understand it, and the degree to which services necessary for health care can be used."

Health information literacy consists of rapid estimate of adult literacy in medicine and

functional literacy. The rapid estimate of adult literacy in medicine (REALM) refers to an understanding of the 66 common terms used in medical institutions related to disease or body. Functional health information literacy refers to the ability to read and understand health-related data, such as medication manuals, appointments, and examination-related instructions, and to take appropriate health actions (4). Previous studies on the relations and effects of the health information literacy on health in Korea include 'a study on the development of a Korean health information literacy evaluation tool' (5), and 'a study on the development of Health Literacy Scale for Korean Teens (KHLS-Teen)' (6).

Overseas studies include 'a research on the relationship between health promotion behavior and health information literacy of Taiwanese teenagers' (7) and 'a study on the development and validity of health information literacy prediction model for Europeans' (8). As identified in these previous studies, low health information literacy becomes a negative factor in proper health-related activities, and furthermore, a barrier of effective communication in doctor-patient resulting relations, in deterioration in the quality of medical services. In this respect, health information literacy is closely related to health promotion.

Recently, with the Fukushima nuclear disaster, interest in radiation safety has been increasing in our lives. Radiation is small particle radiation or electromagnetic wave that causes ionization or excitation. It is important to avoid exposure as much as possible since human exposure to radiation can cause biological damage such as cancer. Like this, despite the importance of radiation related to health, the public's understanding of radiation is not sufficient. As a result, researches on the role of radiologists to improve the health information literacy of patients in relation to radiation therapy have been conducted recently (9). In addition, researches measuring on radiation-related knowledge of the general public are being conducted (10, 11).

These studies are all related to safety from radiation, which is very meaningful for health. However, few studies have been conducted to measure literacy level associated with radiation safety so far. In other words, as most of the literacy measurement tools previously studied in relation to health measure general literacy in the health and medical field, it is not enough to measure the literacy of the radiation field, which has become a hot issue in recent years.

If healthcare providers are informed of the level of literacy associated with the radiation safety of the patient during the course of treatment, they can maximize the effectiveness of the treatment by providing more appropriate terminology or instructions to the patient. In this respect, the development of a tool to measure safety-related literacy radiation necessary and significant. Based on previous studies that the degree of health information literacy is closely related to health promotion activities, it is considered that radiation safety literacy is also closely related to health. However, since there is no measurement tool for this purpose, this study intends to develop a tool for the measurement of radiation safety literacy (MRSL) and to identify actual condition through survey.

In addition, we will evaluate the statistical aspects by analyzing the reliability of the developed measurement tool and checking the validity using correlation coefficient with REALM. We will also examine whether developed measurement tool is a meaningful tool for setting the criteria for literacy level determination using the receiver operating characteristic (ROC) curve. The results of the radiation safety literacy assessments based on the tools developed in this study can be useful for evaluating patients for the appropriate communication between healthcare providers and patients and for evaluating the radiation safety literacy for the public.

MATERIALS AND METHODS

Development procedure of radiation safety literacy measurement tool

The tool development procedure for radiation safety literacy measurement was performed by applying a four-step procedure for

Int. J. Radiat. Res., Vol. 19 No. 1, January 2021

development the of general literacy measurement tools (12). First, the researchers of this study selected terms that had high frequency of occurrence more than 10 times in the books such as radiation-related safety education materials, radioiodine therapy guide, radiation terminology dictionary (13), radiation and life (14), and nuclear energy and radiation (15) as preliminary terms. Second, the content validity of the items extracted in step 1 was evaluated using experts consisting of 1 radiologist working in the dep. of radiology, 2 professors of dep. of radiological science majoring in radiation therapy and 1 angiography nurse. The criteria for the selection of experts were for those who could explain and have enough knowledge about radiation to assist in development of radiation literacy measurement tools during evaluation process of the content validity. Third, another expert group was composed of 2 radiologists, which included one from a department of radiation oncology and another from a department of nuclear medicine, and 1 professor of radiology majoring in magnetic resonance imaging to examine the difficulty level of each item by verifying face validity of the items extracted in step 2. Fourth, a survey must be conducted for statistical assessment of the final terms selected according to the above three steps. At this time, the result of the response is greatly affected by how the response scale is constructed. Therefore, a desirable scale of survey responses was constructed by the consultation of a professor of the Korean Language department. Previous studies measuring medical information comprehension presented only 'yes' or 'no' to check the respondents' understanding of the meaning of the proposed term. However, although this dichotomous response type is expected to have active responds, it not only increases the response burden of respondents but also respondents who know to some extent may respond "no." A response of "so so" may reduce the burden of respondents but may lead to passive response. Therefore, in order to solve these problems, the response category of this study is composed of 4 scale response types (1 point: I don't know-4 points: I know well) on the

advice of the professor of the Korean Language department.

Data collection

The questionnaire presented in table 1 was made and used for data gathering for the statistical assessment of developed radiation safety literacy measurement tool. The survey was conducted from October 1 to October 29, 2018 for students of I university and samples of n=280 were collected by convenience sampling to allow them to respond according to the self-administered method. Demographic information of the sample is shown in table 2. The survey was conducted on students who only voluntarily expressed their willingness participate after explaining the purpose and contents of the study fully before obtaining the response. Respondents were provided with a coffee coupon. Data input and analysis for statistical assessment was done using IBM SPSS 25.

Table 1. Questionnaire for data gathering (part).

How much do you know about the following health care and radiation related terms? ① I don't know ② I do not know well (4) I know well ③ I know a little **REALM** term 1 2 3 4 fat flu llig dose MRSL 1 2 4 term 3 nuclear power generation proton therapy radiation radiation activity

Table 2. Demographics of sample.

| 3 1 | | | | |
|-----------------|----------------------------|--|--|--|
| characteristic | summary value | | | |
| age mean (SD) | 20.89(3.98) | | | |
| male/female (%) | 124(44.3)/156(55.7) | | | |
| Major (%) | radiological science(34.2) | | | |
| | nursing(18.0) | | | |
| | accounting(12.5) | | | |
| | textile(13.6) | | | |
| | engineering(14.0) | | | |
| | sports couching(7.7) | | | |

RESULTS

The result of measurement tool development

A total of 79 terms were selected through the extraction of preliminary terms, the first stage for the development of a radiological safety literacy measurement tool. However, among these, terminology such as linear accelerator and brachytherapy and general terms such as radiation and radon were mixed. One of the main purposes of this study is to measure the radiation safety related literacy of the general Therefore. terms public. 26 including deterministic effect, stochastic effect, thoron, spent nuclear fuel, and radioactive attenuation, which are judged to be too professional or low validated were excluded on the advice of a group of four experts. In addition, some terms have been modified to more general terms, such as 'ultrasound examination ' for 'ultrasound scan ' and 'shielding material' for 'shielding block'.

Terms such as X-rays, computed tomography, and Sievert were written X-ray, CT, and Sv in parentheses in English, respectively. In addition, term of 'electromagnetic waves' was revised into 'electric waves' more commonly used in our lives. For the 53 terms selected through the content validity evaluation, which is the second step, the face validity evaluation for the difficulty evaluation was performed as the third step.

As a result, unusual terms such as 'containment' and 'decontamination' and relatively easy terms such as 'energy' were excluded and 'internal exposure' and 'external exposure' were simply modified to 'radiation exposure'. Finally, a total of 46 terms were selected as shown in table 3.

Table 3. Final extracted radiation safety related terms.

| Table 3. Final extracted radiation safety related terms. | | | | | | |
|--|--|----------------|--|--|--|--|
| Item number | Term | Item number | Term | | | |
| 1 | nuclear power generation | 24 | radioisotope | | | |
| 2 | proton therapy | 25 | beta rays | | | |
| 3 | radiation | 26 | gamma ray | | | |
| 4 | radiation activity | 27 | alpha ray | | | |
| 5 | Radioactive material | 28 | radiation therapy | | | |
| 6 | electromagnetic wave | 29 | radon | | | |
| 7 | X-ray | 30 | uranium | | | |
| 8 | Computed tomography | 31 | radiation exposure | | | |
| 9 | Magnetic resonance imaging | 32 | protection of radiation harm | | | |
| 10 | Ultrasound examination | 33 | shielding material | | | |
| 11 | cosmic radiation | 34 | irradiated food | | | |
| 12 | natural radiation | 35 | Sievert | | | |
| 13 | radioactive iodine | 36 | Radioactive contamination | | | |
| 14 | radioactive waste | 37 | Department of Nuclear Medicine | | | |
| 15 | Chernobyl nuclear accident | 38 | Department of Radiation Oncology | | | |
| 16 | Fukushima nuclear accident | 39 | Department of Radiology | | | |
| 17 | atomic bomb | 40 | angiography | | | |
| 18 | positron emission tomography | 41 | half-life | | | |
| 19 | high level / low level(radioactivity) | 42 | Radioactive decay | | | |
| 20 | radiation sensitivity | 43 | ultraviolet rays | | | |
| 21 | nondestructive inspection | 44 | infrared rays | | | |
| 22 | International Atomic Energy Agency | 45 | visible rays | | | |
| 23 | cancer | 46 | Genetic modification | | | |

Descriptive statistics of 46 MRSL items

According to the survey using n=280 sample for the radiation safety literacy measurement developed in section 3.1, the mean and standard deviation of the response scores for 46 items are shown in table 4. The least understood term was

Int. J. Radiat. Res., Vol. 19 No. 1, January 2021

[DOI: 10.29252/ijrr.19.1.41]

'protection of radiation harm' of 1.80 points, followed by 'radioactive iodine' (1.83 points) and 'Sievert' (1.87 points). On the other hand, most understandable terms 'cancer' (3.35 points) and 'X-ray' (3.28 points), which are commonly encountered in everyday life. Meanwhile, the score of n = 280 for the sum of 46 MSRL items was 116.71 ± 27.38 and the score for the sum of 66 REAL items was 211.92 ± 30.44 .

Table 4. Mean and standard deviation of 46 MRSL items.

| Item number | Mean ± SD | Item number | Mean ± SD |
|-------------|-------------|-------------|--------------|
| 1 | 2.91±0.8130 | 24 | 2.37± (1.055 |
| 2 | 2.04±0.856 | 25 | 2.41±1.027 |
| 3 | 3.15±0.717 | 26 | 2.47±1.035 |
| 4 | 3.14±0.714 | 27 | 2.46± 1.024 |
| 5 | 2.96±0.830 | 28 | 3.03±1.485 |
| 6 | 3.18±1.939 | 29 | 2.12± 1.028 |
| 7 | 3.28±0.660 | 30 | 2.39±0.976 |
| 8 | 2.92±0.879 | 31 | 2.80± 0.961 |
| 9 | 2.89±0.957 | 32 | 1.80±0.890 |
| 10 | 3.12±0.789 | 33 | 1.94±1.039 |
| 11 | 2.09±0.951 | 34 | 1.85±0.932 |
| 12 | 2.16±0.963 | 35 | 1.87±1.053 |
| 13 | 1.83±0.819 | 36 | 2.84±1.095 |
| 14 | 2.48± 0.951 | 37 | 2.14±0.996 |
| 15 | 2.50±1.112 | 38 | 2.10±0.997 |
| 16 | 3.12± 0.795 | 39 | 2.71±0.974 |
| 17 | 3.08±0.774 | 40 | 2.20±1.063 |
| 18 | 2.11±0.980 | 41 | 2.33±1.099 |
| 19 | 1.90±0.934 | 42 | 2.26±1.046 |
| 20 | 2.13±1.018 | 43 | 3.18±0.699 |
| 21 | 1.93± 0.955 | 44 | 3.14±0.762 |
| 22 | 2.24±0.998 | 45 | 2.97± 0.876 |
| 23 | 3.35±0.688 | 46 | 2.96± 0.887 |

DISCUSSION

Health literacy is the level of the individual ability to obtain, process, and understand the basic health information and services necessary to make right health decisions. In other words, health literacy means the ability to obtain, read. understand, and utilize health information so that an individual can make correct judgments regarding his or her medical use. The most widely used and representative test tools for measuring literacy in the health care sector are Rapid Estimate of Adult literacy (REALM) and Test of Functional Health Literacy in Adult (TOFHLA) (16).

Since then, various variant tools of REALM and TOFHLA have been developed and used. Some examples include Rapid Estimate of Adult Literacy in Medicine, Revised (REALM-R), Rapid Estimate of Adult Literacy in Medicine—Short Form, and Short Assessment of Health Literacy for Spanish Adults (SAHLSA-50) (17).

However, these health literacy measurement tools are all for measuring general health-related literacy. Meanwhile, after 2011 Fukushima nuclear disaster, radiation safety has emerged as an important issue related to health for neighboring countries in East Asia, including Japan. High literacy level related to radiation safety increase the ability to acquire relevant information, resulting in acquiring high-quality medical services. In other words, the higher the radiation safety literacy, the safer it is from radiation. For this reason, it is necessary to identify the actual condition of radiation safety literacy and to carry out related research to protect our health from radiation. Accordingly, many researches related to radiation safety have been conducted recently. Recent studies related to radiation safety include 'a study suggesting the need to provide appropriate educational materials to parents to assist in decision making during computed tomography for children'(18, 19), 'a study of a pediatric emergency department showing that the lower the health literacy of the guardian, the lower the degree of radiographic examination' (20), and 'a study of the relationship between radiation concern and health literacy among residents living in evacuation area and non-evacuation area after the Fukushima nuclear accident' (21). However, there are few measurement tools that can measure the radiation safety literacy of the public in the previous studies related to radiation safety. Therefore, this study developed 46 MRSL in accordance with the four-step procedure outlined in figure 1. However, it is difficult to discuss the relative superiority of MRSL as there is no appropriate previous study to compare with the developed measurement tool, MRSL. In

Choi and Cho / Development of a radiation literacy measurement tool

this regard, this study will discuss the adequacy of the developed radiation safety literacy measurement tool through the various statistical aspects of MRSL – reliability, validity, and the discussion on whether a measurement tool developed using receiver operating characteristic (ROC) curve is meaningful for setting criteria for literacy level determination.

First, reliability analysis was conducted to find out how consistent the measurement tool developed through this study showed. The reliability of this study was determined using Cronbach's "\alpha" coefficient since the response of the study was the multiple-choice scale (1 point: I don't know ~ 4 points: I know well) compared to response of dichotomous, which use KR-20 confidence coefficient. The Cronbach's coefficient, an internal consistency measure for all 46 items, was very high at α =0.963, indicating that the reliability of the radiation safety literacy measurement tool of table 3 is considered to be very high. Meanwhile, when the specific items were removed, only two items of the 6th (electronic wave) and 28th (radiation therapy) had Cronbach's "α" coefficient greater than 0.963. However, the overall reliability coefficient value did not increase significantly to 0.966 and 0.964 respectively when these items were removed. Consequently, these two items were not removed from the radiation safety literacy measurement tool. As the Cronbach's "α" coefficient for the rapid estimate of adult literacy in medicine (REALM) of 66 items presented in the previous study was 0.977, the reliability coefficient of MRSL was as high as the previous studies.

Next, in order to evaluate the criterion validation of the radiation safety literacy measurement tool developed in this study, the correlation coefficient with REALM, which was proved in previous studies, was examined. To this end, the sum of 66 items (66 to 264 points) for measuring REALM and the sum of 46 items (44 to 176 points) developed in this study to measure radiation safety literacy (MRSL) were obtained. Pearson's correlation coefficient was found to be 0.448, showing statistically significant (p<0.01) at the significant level of 1%. Therefore, the measurement tool developed

to measure radiation safety literacy in this study was evaluated to be statistically valid. As another method to assess the validity, response distribution of the radiation safety literacy measurement tool developed in this study (figure 2) was examined whether it follows the normal distribution. The result of the Kolmogorov-Smirnov normality test showed that it was suitable for the normal distribution at the significant level of 5% with p=0.051.




Figure 1. Flowchart of radiation safety literacy measurement tool development.




Figure 2. Response distribution of MRSL score.

Finally, the ROC curve shown in figure 3 is used to help determine whether the respondent's literacy level is high or low based on the MRSL ⁽²²⁾. In this curve, the horizontal

Int. J. Radiat. Res., Vol. 19 No. 1, January 2021

axis represents 1-specificity and the vertical axis represents sensitivity. As the point is upward and on the left on the graph, the sensitivity is which means greater. more accurate determination. In table 5, sensitivity refers to the probability (A/(A+C)) that accurately detects low MRSL in person with low radiation safety literacy level, and specificity means the probability (D/(B+D)) to determine that a person with a high radiation safety literacy has high level of MRSL. In figure 3, the area below the MRSL curve is 0.709, with a significant probability of p<0.01, indicating that the judgment method using MRSL is a statistically valuable (23).

Table 5. Result for assessment of literacy level.

| MRSL | REALM | | |
|--------|-------|------|--|
| IVIKSL | low | high | |
| low | Α | В | |
| high | С | D | |




Figure 3. ROC for decision criterion making.

CONCLUSION

The MRSL tool consisting of 46 terms developed in this study has been proven its reliability and validation in statistical terms. In addition, from the test on the ROC curve, the radiation safety literacy level evaluation using MRSL was found to be a statistically valuable

Int. J. Radiat. Res., Vol. 19 No. 1, January 2021

judgment method. Therefore, it is believed that the radiation safety literacy measurement tool developed in this study can be very useful for conducting future researches in East Asia affected by the Fukushima nuclear disaster. Furthermore, the results of the radiation safety literacy assessment measured based on MRSL can be used for appropriate communication between health care providers and patients, or for developing and providing educational materials by level for the public.

Conflicts of interest: Declared none.

REFERENCES

- Davis TC, Long SW, Jackson RH, Mayeauk EJ, George RB, Murhpy PW, Crouch MA (1993) Rapid estimate of adult literacy in medicine: A shortened screening instrument. *Journal of Family Medicine*, 25(6): 391-395.
- Kim M (2018) Development of diabetes education material for older people with low health literacy in rural community, Ph.D. thesis, Mokpo National University, Jeollanam-do.
- 3. Tappe MS and Galer-Unti, RA (2001) Health educators' role in promoting health literacy and advocacy for the 21st century. *Journal of School Health*, **71(10)**: 477-482.
- 4. Hwang TY (2010) Understanding health literacy: Implication for medicine and public health. *Academy Press, p. 5*.
- 5. Kim SS, Kim SH, Lee SY (2005) Health literacy: Development of a Korean health literacy assessment tool. *Journal of Korean Society for Health Education and Promotion*, **22(4)**: 215-227.
- 6. Jang B (2017) Development of Korean adolescent health literacy scale(KHLS-Teen), Ph.D. thesis. *Busan National University*, p. 92.
- 7. Chang LC (2011) Health literacy, self-reported status abd health promoting behaviors for adolescents in Taiwan. *Journal of Clinical Nursing*, **20**: 190-196.
- 8. van der Heide I, Uiters E, Boshuizen H, Rademakers J (2015) Health literacy in Europe: The development and validation of health literacy prediction models. *European Journal of Public Health*, **26(6)**: 906-911.
- Montgomery L (2015) Supporting radiation therapy patients with limited health literacy. Journal of Medical Imaging and Radiation Sciences, 46: 102-107.
- Kim SH and Cho K (2016) Analysis of university student awareness of radiation exposures from consumer products. *Journal of Radiation Protection*, 41(1): 57-70.
- 11. Geofery L, Basirat M, Eze CU, Chigozie NL, Auwal A, Kalu O, Bobuin NF, Mohammed N, Moi AS, Mathew A G (2015) Evaluation of the knowledge of non-ionizing radiation among final year students of colledge of medical science

Choi and Cho / Development of a radiation literacy measurement tool

- university of Maiduguri. Int Res J Pure and Applied Physics, 3(3): 8-14.
- Kim SH, Jung S, Park K, Lee SH, Choi Y, Lee WH, Choi KH (2016) Development of the Korean screening tool for anxiety disorders: Review of current anxiety scales and development of preliminary item pools. *Korean Journal of Clinical Psychology*, 35(3): 630-644.
- 13. Koh SK (2010) The radiation dictionary. Hyunmoonsa, Seoul, Korea.
- 14. Korea Academy of Nuclear Safety (2015) Radiation and life, Korea Academy of Nuclear Safety, Seoul, Korea.
- Korea Academy of Nuclear Safety (2015) Nuclear energy and radiation, Korea Academy of Nuclear Safety, Seoul, Korea.
- 16. Hwang TY (2010) Understanding heal literacy: Implication for medicine and public health, Academy Press, Seoul.
- Bass PF, Wilson JF, Griffith CH (2003) A shortened instrument for literacy screening. *Journal of General Internal Medicine*, 18(12): 1036-1038.
- 18. Goske MJ and Bulas D (2009) Improving health literacy:

- Informed decision-making rather than informed consent for CT scans in children. *Pediatric Radiology*, **39**: 901-903.
- Bulas D, Goske MJ, Applegate K, Wood B (2009) Image gentry: Improving health literacy for parents about CT scans for children. *Pediatric Radiology*, 39: 112-116.
- Morrison AK, Brousseau DC, Brazausks R, Levas MN (2015)
 Health literacy affects likelihood of radiaology testing in
 the pediatric emergency department. The Journal of Pediatrics, 166(4): 1037-1041.
- 21. Kuroda Y, Iwasa H, Orui M, Moriyama N, Nakayama C, Yasumura S (2018) Association between health literacy and radiation anxiety among residents after a nuclear accident: Comparison between evacuated and non-evacuated area. *Int J Environ Res and Public Health*, 15: 1463-1475.
- Lisa DC, Katharine AB, Edward JB (2004) Brief questions to identify patients with inadequate health literacy. Family Medicine, 36(8): 588-594.
- Park MR, Lee J, Cho J (2016). Statistical analysis of medical data using SPSS. Freedom-Academy, 275-276, Seoul, Korea.