

## Evaluation of activity concentration of natural radionuclides and lifetime cancer risk in soil samples at two tertiary institutions in Owerri, Imo State, Nigeria

B.C. Eke<sup>1</sup>, U.M. Ukewuihe<sup>1</sup>, I.R. Akomolafe<sup>2\*</sup>

<sup>1</sup>Department of Physics, School of Physical Sciences, Federal University of Technology, Owerri, Nigeria

<sup>2</sup>Department of Physical Sciences, Redeemer's University, Ede, Nigeria

### ABSTRACT

#### ► Original article

##### \*Corresponding author:

Idowu Richard Akomolafe, Ph.D.,  
E-mail:  
[idowurakomolafe@gmail.com](mailto:idowurakomolafe@gmail.com)

Received: May 2021

Final revised: November 2021

Accepted: December 2021

*Int. J. Radiat. Res.*, July 2022;  
20(3): 671-678

DOI: 10.52547/ijrr.20.3.22

**Keywords:** Radiological risks, natural radioactivity, soil samples, Nigeria.

**Background:** Environmental radioactivity measurement of soil samples from the densely populated community has become imperative considering the radiological exposure associated with primordial radionuclides. **Materials and Method:** Sixty soil samples were collected at different locations within Alvan Ikoku Federal College of Education (AIFCE) and Federal Polytechnic Nekede (FPN), Owerri. Radioactivity measurements were carried out by the method of gamma-ray spectrometry with thallium doped sodium iodide [NaI(Tl)] detector. **Results:** The mean activity concentrations of  $88.41 \pm 1.51$ ,  $20.69 \pm 3.56$  and  $25.04 \pm 0.71$  Bqkg<sup>-1</sup> for  $^{40}\text{K}$ ,  $^{226}\text{Ra}$  and  $^{232}\text{Th}$ , respectively, were obtained in soil samples of AIFCE, while  $92.97 \pm 1.50$ ,  $20.48 \pm 3.21$  and  $22.36 \pm 0.70$  Bqkg<sup>-1</sup> for  $^{40}\text{K}$ ,  $^{226}\text{Ra}$  and  $^{232}\text{Th}$ , respectively, were obtained in soil samples of FPN. These values are below the average world value of 420, 32 and 45 Bqkg<sup>-1</sup> for  $^{40}\text{K}$ ,  $^{226}\text{Ra}$  and  $^{232}\text{Th}$ , respectively, as recorded in UNSCEAR 2000 report. The calculated absorbed dose, annual effective dose and other radiological hazard indices were below the recommended safe limit. The mean calculated values of the excess lifetime cancer risk for AIFCE and FPN are  $0.12 \pm 0.03$  and  $0.11 \pm 0.03$ , respectively. **Conclusion:** Based on the results from the present study, it is evidence that the activity concentrations and other radiological parameters are within the world's safe limit, indicating that soil samples from the AIFCE and FPN, Owerri are free from radioactive contamination and do not pose a threat to the two communities.

### INTRODUCTION

Human beings are constantly exposed to ionizing radiation from natural sources that have become an unavoidable part of life on earth<sup>(1)</sup>. Human exposure to terrestrial radiation is primarily by gamma radiation from radionuclides in the  $^{238}\text{U}$  and  $^{232}\text{Th}$  series and non-decay  $^{40}\text{K}$ . This exposure exceeds that from all artificial sources combined for most people<sup>(1)</sup>. Natural radioactivity can be found in various geological formations, including the earth's crust, rocks, soils, plants, water, and air. Geological and geographical conditions primarily determine the natural radioactive concentration, and it can be found at various levels in soils from different geological regions<sup>(1,2)</sup>.

The radionuclide activity concentration in soil samples is one major determinant in assessing the natural background radiation<sup>(2)</sup>. Radionuclides are delivered to the soil by rain and flows as rocks disintegrate through the activity of weathering<sup>(3)</sup>. The report has shown that various radiation levels exist with different types of rocks. For instance, igneous rocks are usually associated with higher radiation levels and sedimentary rocks with lower

radiation levels<sup>(1)</sup>.

The relationship between radiation exposure and cancer cases cannot be downplayed, as reports from epidemiological studies indicate a dose-response hypothesis<sup>(4)</sup>. This hypothesis suggests any rise in radiation dose could yield an increase in cancer risk even at a small dose. A linear, no-threshold relationship exists between radiation dose and cancer occurrence<sup>(4,5)</sup>. That is, the dose of radiation receives by an individual from ionizing radiation tends to initiate cancer. The parameter commonly used to assess the impact of cancer on the population study is excess lifetime cancer risk which is the probability that an individual will develop cancer over their lifetime of exposure<sup>(4,5)</sup>.

Radioactivity measurement of environmental samples remains one significant way to determine the level of natural radionuclide around us and ascertain their level of hazard to man<sup>(6)</sup>. Studies have shown that soil is a continuous source of radiation exposure and acts as a medium of migration to transfer radionuclides to biological systems<sup>(7,8)</sup>.

Alvan Ikoku Federal College of Education (AIFCE) is one of the tertiary institutions in Imo State, Nigeria. The defunct Eastern Nigeria Government established

the Institution in April 1963. The Federal Polytechnic Nekede (FPN) was created initially as Government Technical College by the Imo State government in 1978 but later metamorphosed into Federal Polytechnic Nekede by the Federal Government of Nigeria on the 7<sup>th</sup> April 1993.

Environmental radioactivity measurement of soil samples from the densely populated community cannot be downplayed as such measures can be used to estimate the level of radioactive contamination of the environment. In addition, it can serve as a baseline for the epidemiological study of the community in case of any discharge of radioactive material to the environment. Therefore, it becomes imperative to consider the area with a dense population and ascertain their radiation exposure level. Furthermore, to the best of our knowledge, no study has been conducted to determine soil samples' radioactivity level from AIFCE and FPN, Owerri.

Thus, the study aimed to evaluate the activity concentration of natural radionuclides ( $^{226}\text{Ra}$ ,  $^{232}\text{Th}$ , and  $^{40}\text{K}$ ) in soil samples from AIFCE and FPN. In addition, the study calculated the radiological hazard indices and the excess lifetime cancer risk associated with the soil samples from these institutions.

## MATERIALS AND METHODS

### Samples collection and preparation

Sixty soil samples were collected randomly from different locations within the campuses; thirty soil samples from AIFCE and the remaining soil samples from FPN. The samples were collected to a depth of 150 mm below the surface, placed in polythene bags, and carefully labelled (9). The maps of the sample locations are shown in figures 1 and 2. After that, the collected soil samples were transferred to the Radiation and Health Physics Research Laboratory at the Department of Physics, University of Ibadan, Nigeria, where they were prepared for radioactivity counting. Briefly, the soil samples were oven-dried at 110°C to remove the moisture until a constant weight was attained. After which, the soil samples were grounded, homogenized and sieved with a 2.0 mm mesh sieve (9). The sieved soil samples weighing 0.2kg were packed into cleaned airtight plastic containers of uniform size and sealed. The sealed samples were stored in a dried place and left for a minimum of 28 days to ensure radioactive secular equilibrium between  $^{226}\text{Ra}$ ,  $^{232}\text{Th}$  and their short-lived daughter products (10).

### Measuring System

Radioactivity measurement of soil samples was carried out by the method of gamma-ray spectrometry using a lead-shielded 76 mm × 76 mm thallium doped sodium iodide NaI (Tl) detector. The detector was connected with a Canberra Series 10+

Multichannel Analyser (MCA) via a preamplifier. The MCA is a comprehensive system that includes all spectroscopic analysis operations. The spectrometer has a resolution of 8% efficiency at an energy of 0.662 MeV ( $^{137}\text{Cs}$ ), capable of differentiating the radionuclides used for the measurement. The photo-peak energy of 1.460 MeV was used to identify  $^{40}\text{K}$ , 1.760 MeV for  $^{226}\text{Ra}$  ( $^{238}\text{U}$ ) and 2.614 MeV for  $^{232}\text{Th}$ . A standard reference soil sample from Rocketdyne Laboratories, California, USA, was used for the efficiency calibration. The reference sample is traceable to a mixed standard gamma source (Ref No 48722-356) by Analytic Inc., Atlanta, GA, USA. The reference sample was placed on top of the detector and counted for 10 hours (36000s). By removing counts attributable to Compton scattering of higher peaks and other background sources from the peaks' total area, the net area under the corresponding peaks in the energy spectrum was determined. Each sealed sample was placed on top of the detector and counted for the same time as the reference sample based on the stored spectra.




Figure 1. Sample locations at the Alvan Ikoku Federal College of Education, Owerri.




Figure 2. Sample locations at the Federal Polytechnic Nekede, Owerri.

### Activity concentration and radiological parameters

The activity concentrations of the radionuclides in the soil samples were calculated using equation (1) (10, 11).

$$C(\text{Bqkg}^{-1}) = kC_n \quad (1)$$

where  $C_n$  is the count rate under the corresponding peak,  $k = \frac{1}{\epsilon P_\gamma M_s}$ ,  $P_\gamma$  is the absolute

transition probability of the specific gamma-ray,  $C$  is the activity concentration of the radionuclide of soil samples given in  $\text{Bqkg}^{-1}$ ,  $\varepsilon$  is the detector efficiency at the specific gamma-ray energy,  $t$  is the counting time in seconds and  $M_s$  is the mass of the sample (kg). The gamma-ray detector's detection limit (DL) defines its operational capability without the sample's influence (11). This calculation was performed using equation (2).

$$DL (\text{Bqkg}^{-1}) = 4.65 \frac{(C_b)^{1/2}}{t_b} k \quad (2)$$

Where;  $t_b$  is the background counting time in second,  $C_b$  is the net background count in the corresponding peak,  $k$  is the conversion factor given in equation (1). The present study's measurement system showed that soil samples' detection limits were 16.96, 3.65 and 4.43  $\text{Bqkg}^{-1}$  for  $^{40}\text{K}$ ,  $^{226}\text{Ra}$  and  $^{232}\text{Th}$ , respectively. Any activity concentration values below these numbers were taken below the detector's detection limit (BDL).

#### Absorbed dose rate

The absorbed dose rate in the air from exposure to natural primordial radionuclides was calculated using the activity concentration of radionuclides results. The absorbed dose rate ( $D (\text{nGyh}^{-1})$ ) in the air helps quantify the amount of radiation absorbed by a body at 1 m above the ground due to  $^{40}\text{K}$ ,  $^{226}\text{Ra}$  and  $^{232}\text{Th}$ . The absorbed dose rate in the air was calculated using equation (3) as given in the UNSCEAR report (1, 10).

$$D(\text{nGyh}^{-1}) = 0.462C_{\text{Ra}} + 0.604C_{\text{Th}} + 0.0417C_{\text{K}} \quad (3)$$

Where;  $C_{\text{Ra}}$ ,  $C_{\text{Th}}$  and  $C_{\text{K}}$  are the activity concentration in  $\text{Bqkg}^{-1}$  of  $^{226}\text{Ra}$ ,  $^{232}\text{Th}$  and  $^{40}\text{K}$ , respectively.

#### Annual effective dose (AED)

To assess the annual effective dose by a member of the community, we considered two factors. Firstly, the absorbed dose rate in  $\text{nGy}^{-1}$  was converted to human effective dose  $\text{Sv}^{-1}$  using the conversion factor of 0.7  $\text{SvGy}^{-1}$ . Secondly, the average time an individual is exposed to outdoor or indoor radiation was put into consideration. The United Nations Scientific Committee on the Effects of Atomic Radiation (1) recommended 0.2 and 0.8 for outdoor and indoor occupancy factors, respectively. This study considered only outdoor exposures from gamma-ray sources due to the concentrations of naturally occurring radioactive materials (NORMs). The annual effective dose resulting from the absorbed dose rate values was calculated using equation (4) (1).

$$AED = D (\text{nGy}^{-1}) \times 8760 \text{hy}^{-1} \times 0.2 \times 0.7 \text{SvGy}^{-1} \quad (4)$$

Where;  $AED$  is the annual effective dose ( $\mu\text{Sv}^{-1}$ ),

$D$  is the absorbed dose rate in the air.

#### Radium equivalent activity ( $\text{Ra}_{\text{eq}}$ )

Radium equivalent is a regular radiological index often used to compare the specific activities of soil samples containing  $^{40}\text{K}$ ,  $^{226}\text{Ra}$ , and  $^{232}\text{Th}$ , radionuclides by a single quantity, which considers the radiation hazards associated with them (10, 12-15).  $\text{Ra}_{\text{eq}}$  was calculated using equation (5):

$$\text{Ra}_{\text{eq}} (\text{Bqkg}^{-1}) = C_{\text{Ra}} + 1.43C_{\text{Th}} + 0.077C_{\text{K}} \quad (5)$$

Where  $C_{\text{Ra}}$ ,  $C_{\text{Th}}$  and  $C_{\text{K}}$  are as defined in equation (3). Radiation hazard from soil samples can only be negligible if the value of  $\text{Ra}_{\text{eq}}$  is less than 370  $\text{Bqkg}^{-1}$ .

#### External and internal hazard index

External hazard index ( $H_{\text{ex}}$ ) and internal hazard index ( $H_{\text{in}}$ ) are essential criteria used to measure the level of exposure to radon ( $^{222}\text{Rn}$ ), a daughter of  $^{226}\text{Ra}$  in the  $^{238}\text{U}$  decay series. Radon has been known to be dangerous to respiratory organs if inhalation takes place. This radionuclide and its radioactive progenies pose severe internal exposure. Equations (6) and (7) were used to calculate external and internal hazard indices.

$$H_{\text{ex}} = \frac{C_{\text{Ra}}}{370 \text{Bqkg}^{-1}} + \frac{C_{\text{Th}}}{259 \text{Bqkg}^{-1}} + \frac{C_{\text{K}}}{4810 \text{Bqkg}^{-1}} \quad (6)$$

$$H_{\text{in}} = \frac{C_{\text{Ra}}}{185 \text{Bqkg}^{-1}} + \frac{C_{\text{Th}}}{259 \text{Bqkg}^{-1}} + \frac{C_{\text{K}}}{4810 \text{Bqkg}^{-1}} \quad (7)$$

Where  $C_{\text{Ra}}$ ,  $C_{\text{Th}}$  and  $C_{\text{K}}$  are as defined in equation (3). Radiation hazard from soil samples is considered negligible if the value of both indices is less than unity (1).

#### Annual Gonadal Dose Equivalent (AGDE)

The annual gonadal dose equivalent considers the radiation dose received by the reproductive organs, bone marrows and bone cells due to activity concentrations of natural radionuclides in soil samples. The purpose is to determine the extent of radiation hazard to these organs, which happens to be the body's highly radiosensitive organs. Research has shown that an increase in AGDE could damage the bone marrow, causing the destruction of red blood and white blood cells replenished. This condition causes leukaemia, a type of blood cancer that is lethal. AGDE was estimated using equation (8) (4, 16, 17).

$$AGDE (\mu\text{Sv}^{-1}) = 3.09C_{\text{Ra}} + 4.18C_{\text{Th}} + 0.314C_{\text{K}} \quad (8)$$

Where;  $C_{\text{Ra}}$ ,  $C_{\text{Th}}$  and  $C_{\text{K}}$  are the activity concentrations of radium, thorium and potassium, respectively in  $\text{Bqkg}^{-1}$ , 3.09, 4.18 and 0.314 are conversion factors.

#### Excess Lifetime Cancer Risk (ELCR)

Excess lifetime cancer risk is a parameter that

measures the probability of cancer risk to any population due to radiation exposure. It is expressed as a number representing the number of additional cancers expected in a given number of people exposed to a carcinogen at a specific dose. It was calculated based on the estimated annual effective dose. Equation 9 was used to calculate ELCR as provided by International Commission on Radiological Protection (ICRP) publication 60 (3, 4, 17-19).

$$\text{ELCR} = (\text{AED} \times \text{DL} \times \text{RF}) \times 10^{-3} \quad (9)$$

Where; AED is the annual effective dose, DL is the average duration life span (taken as 70 years), and RF is the risk factor given as  $0.05\text{Sv}^{-1}$  that is a fatal cancer risk for stochastic effects in any given population based on ICRP.

### Statistical analysis

Statistical package for social science (SPSS 20.0) was used for the statistical analysis of data. Results are reported as means  $\pm$  SD (standard deviation).

## RESULTS

The results of the activity concentration of the natural radionuclides as well as absorbed dose and annual effective dose in measurements of the soil samples from AIFCE are presented in table 1. The activity concentrations varied from  $27.85 \pm 1.28 \text{ Bqkg}^{-1}$  to  $122.10 \pm 1.68 \text{ Bqkg}^{-1}$  for  $^{40}\text{K}$  with a mean value of  $88.41 \pm 1.51 \text{ Bqkg}^{-1}$ , whereas for  $^{226}\text{Ra}$ , it ranged from BDL to  $32.74 \pm 3.92 \text{ Bqkg}^{-1}$  with a mean value of  $20.69 \pm 3.56 \text{ Bqkg}^{-1}$ , and for  $^{232}\text{Th}$ , it varied from BDL to  $39.10 \pm 0.54 \text{ Bqkg}^{-1}$  with a mean value of  $25.04 \pm 0.71 \text{ Bqkg}^{-1}$ .

The activity concentration values for soil samples at FPN ranged from  $30.90 \pm 1.47$  to  $147.18 \pm 1.51 \text{ Bqkg}^{-1}$  with a mean value of  $92.97 \pm 1.50 \text{ Bqkg}^{-1}$ , BDL to  $31.11 \pm 4.28 \text{ Bqkg}^{-1}$  with a mean value of  $20.48 \pm 3.21 \text{ Bqkg}^{-1}$ , BDL to  $37.15 \pm 0.81 \text{ Bqkg}^{-1}$  with a mean value of  $22.36 \pm 0.70 \text{ Bqkg}^{-1}$  for  $^{40}\text{K}$ ,  $^{226}\text{Ra}$  and  $^{232}\text{Th}$  respectively (table 3).

Table 1 presents absorbed dose rate result, which ranged from  $7.31$  to  $38.72 \text{ nGyh}^{-1}$  with a mean value of  $28.75 \pm 10.39 \text{ nGyh}^{-1}$  and the annual effective dose that varied from  $8.97$  to  $47.52 \mu\text{Sv}^{-1}$  with a mean value of  $35.28 \pm 12.74 \mu\text{Sv}^{-1}$  in soil samples of AIFCE. Similarly, the absorbed dose rate and the annual effective dose of soil samples in FPN are presented in table 3. The absorbed dose value ranged from  $14.49$  to  $38.24 \text{ nGyh}^{-1}$  with a mean value of  $26.27 \pm 5.99 \text{ nGyh}^{-1}$  and the annual effective dose value varied from  $17.79$  to  $46.92 \mu\text{Sv}^{-1}$  with a mean value of  $32.24 \pm 7.35 \mu\text{Sv}^{-1}$ .

Tables 2 and 4 present the average radium equivalent results in soil samples as  $59.79 \pm 16.98 \text{ Bqkg}^{-1}$  and  $56.78 \pm 13.12 \text{ Bqkg}^{-1}$  for AIFCE and FPN,

respectively. The external radiation hazard index of soil samples in AIFCE varied from  $0.04$  to  $0.23$  with a mean value of  $0.16 \pm 0.05$ , and that of internal radiation hazard index varied from  $0.05$  to  $0.31$  with a mean value of  $0.22 \pm 0.06$ . Moreover, external and internal radiation indices of soil samples in FPN ranged from  $0.08$  to  $0.22$  and  $0.09$  to  $0.30$ , with mean values of  $0.15 \pm 0.04$  and  $0.21 \pm 0.05$ , respectively.

The average annual gonadal dose equivalent and average excess lifetime cancer risk due to activity concentrations of natural radionuclides in soil samples at AIFCE is  $186.00 \pm 50.65 \mu\text{Sv}^{-1}$  and  $0.12 \pm 0.03$ , respectively. At the same time, the average annual gonadal dose equivalent and average excess lifetime cancer risk due to radiation exposure at FPN is  $177.47 \pm 39.20 \mu\text{Sv}^{-1}$  and  $0.11 \pm 0.03$ , respectively.

**Table 1.** Activity concentration of naturally occurring radionuclides, absorbed dose and annual effective dose in soil samples from Alvan Ikoku Federal College of Education.

| Sample ID                     | $^{40}\text{K}$ ( $\text{Bqkg}^{-1}$ ) | $^{226}\text{Ra}$ ( $\text{Bqkg}^{-1}$ ) | $^{232}\text{Th}$ ( $\text{Bqkg}^{-1}$ ) | Absorbed Dose ( $\text{nGyh}^{-1}$ ) | Effective Dose ( $\mu\text{Sv}^{-1}$ ) |
|-------------------------------|--|--|--|--------------------------------------|--|
| AIFCE 1                       | $108.14 \pm 1.21$                      | $15.10 \pm 4.10$                         | $27.40 \pm 0.77$                         | 29.24                                | 35.89                                  |
| AIFCE 2                       | $69.11 \pm 1.43$                       | $23.11 \pm 2.11$                         | $39.10 \pm 0.54$                         | 38.72                                | 47.52                                  |
| AIFCE 3                       | $75.44 \pm 1.59$                       | $14.90 \pm 3.44$                         | $23.45 \pm 0.73$                         | 25.13                                | 30.84                                  |
| AIFCE 4                       | $79.24 \pm 1.61$                       | $9.61 \pm 4.10$                          | BDL                                      | 10.42                                | 12.79                                  |
| AIFCE 5                       | $86.58 \pm 1.70$                       | $16.44 \pm 4.24$                         | $30.15 \pm 0.61$                         | 30.7                                 | 37.68                                  |
| AIFCE 6                       | $100.10 \pm 1.34$                      | $21.22 \pm 4.15$                         | $25.74 \pm 0.73$                         | 30.41                                | 37.32                                  |
| AIFCE 7                       | $111.25 \pm 1.49$                      | $15.74 \pm 4.37$                         | $5.12 \pm 0.84$                          | 14.89                                | 18.27                                  |
| AIFCE 8                       | $86.15 \pm 1.43$                       | $8.62 \pm 2.07$                          | $15.11 \pm 0.74$                         | 17.39                                | 21.34                                  |
| AIFCE 9                       | $75.34 \pm 1.52$                       | $13.10 \pm 2.19$                         | $24.20 \pm 0.88$                         | 24.85                                | 30.49                                  |
| AIFCE 10                      | $93.30 \pm 1.61$                       | $25.54 \pm 2.54$                         | $21.73 \pm 0.72$                         | 29.3                                 | 35.96                                  |
| AIFCE 11                      | $97.94 \pm 1.70$                       | $20.17 \pm 3.78$                         | $25.77 \pm 0.64$                         | 29.88                                | 36.67                                  |
| AIFCE 12                      | $92.23 \pm 1.63$                       | $31.15 \pm 4.11$                         | $29.64 \pm 0.53$                         | 36.88                                | 45.26                                  |
| AIFCE 13                      | $110.44 \pm 1.55$                      | $25.50 \pm 4.27$                         | $30.11 \pm 0.61$                         | 35.57                                | 43.65                                  |
| AIFCE 14                      | $98.45 \pm 1.37$                       | $30.10 \pm 3.10$                         | $27.45 \pm 0.72$                         | 35.26                                | 43.27                                  |
| AIFCE 15                      | $53.22 \pm 1.48$                       | $16.17 \pm 2.55$                         | $29.12 \pm 0.80$                         | 28.47                                | 34.94                                  |
| AIFCE 16                      | $98.15 \pm 1.33$                       | $18.44 \pm 3.98$                         | $19.87 \pm 0.72$                         | 25.25                                | 30.99                                  |
| AIFCE 17                      | $102.25 \pm 1.41$                      | $17.30 \pm 3.74$                         | BDL                                      | 14.43                                | 17.71                                  |
| AIFCE 18                      | $92.40 \pm 1.57$                       | $28.10 \pm 3.01$                         | $28.12 \pm 0.66$                         | 34.59                                | 42.45                                  |
| AIFCE 19                      | $107.77 \pm 1.64$                      | $21.83 \pm 3.98$                         | $23.81 \pm 0.71$                         | 29.72                                | 36.47                                  |
| AIFCE 20                      | $87.10 \pm 1.68$                       | $14.51 \pm 4.17$                         | $25.93 \pm 0.73$                         | 27.11                                | 33.27                                  |
| AIFCE 21                      | $121.24 \pm 1.59$                      | $23.66 \pm 4.28$                         | $21.91 \pm 0.81$                         | 29.82                                | 36.59                                  |
| AIFCE 22                      | $27.85 \pm 1.28$                       | $25.37 \pm 3.17$                         | $20.69 \pm 0.59$                         | 25.73                                | 31.58                                  |
| AIFCE 23                      | $79.41 \pm 1.31$                       | $18.33 \pm 2.18$                         | $27.10 \pm 0.64$                         | 29.18                                | 35.81                                  |
| AIFCE 24                      | $99.12 \pm 1.48$                       | $27.68 \pm 4.44$                         | $23.74 \pm 0.77$                         | 31.79                                | 39.01                                  |
| AIFCE 25                      | $73.15 \pm 1.51$                       | $17.82 \pm 4.78$                         | $29.11 \pm 0.79$                         | 30.03                                | 36.85                                  |
| AIFCE 26                      | $51.78 \pm 1.59$                       | $32.74 \pm 3.92$                         | $30.97 \pm 0.81$                         | 36.71                                | 45.05                                  |
| AIFCE 27                      | $87.23 \pm 1.67$                       | $25.10 \pm 4.10$                         | $31.17 \pm 0.64$                         | 35.1                                 | 43.08                                  |
| AIFCE 28                      | $100.25 \pm 1.38$                      | $23.58 \pm 2.36$                         | $14.12 \pm 0.72$                         | 23.73                                | 29.12                                  |
| AIFCE 29                      | $65.55 \pm 1.53$                       | BDL                                      | BDL                                      | 6.3                                  | 7.73                                   |
| AIFCE 30                      | $122.10 \pm 1.68$                      | $19.14 \pm 4.15$                         | $25.48 \pm 0.69$                         | 65.87                                | 80.83                                  |
| <b>Minimum</b>                | <b><math>27.85 \pm 1.28</math></b>     | <b>BDL</b>                               | <b>BDL</b>                               | <b>7.31</b>                          | <b>8.97</b>                            |
| <b>Maximum</b>                | <b><math>122.10 \pm 1.68</math></b>    | <b><math>32.74 \pm 3.92</math></b>       | <b><math>39.10 \pm 0.54</math></b>       | <b>38.72</b>                         | <b>47.52</b>                           |
| <b>Mean<math>\pm</math>SD</b> | <b><math>88.41 \pm 1.51</math></b>     | <b><math>20.69 \pm 3.56</math></b>       | <b><math>25.04 \pm 0.71</math></b>       | <b><math>28.75 \pm 10.39</math></b>  | <b><math>35.28 \pm 12.74</math></b>    |

**Table 2.** Radium equivalent, hazard indices, annual gonadal dose equivalent and excess lifetime cancer risk in soil samples from Alvan Ikoku Federal College of Education.

| Sample ID      | Ra(Bqkg <sup>-1</sup> ) | H <sub>in</sub>  | H <sub>ex</sub>  | AGDE(μSv <sup>-1</sup> ) | ELCR             |
|----------------|-------------------------|------------------|------------------|--------------------------|------------------|
| AIFCE 1        | 62.61                   | 0.21             | 0.17             | 195.15                   | 0.13             |
| AIFCE 2        | 84.34                   | 0.29             | 0.23             | 256.55                   | 0.17             |
| AIFCE 3        | 54.24                   | 0.19             | 0.15             | 167.75                   | 0.11             |
| AIFCE 4        | 22.05                   | 0.09             | 0.06             | 73.09                    | 0.04             |
| AIFCE 5        | 66.22                   | 0.22             | 0.18             | 204.01                   | 0.13             |
| AIFCE 6        | 65.74                   | 0.23             | 0.18             | 204.59                   | 0.13             |
| AIFCE 7        | 31.63                   | 0.13             | 0.09             | 104.97                   | 0.06             |
| AIFCE 8        | 36.86                   | 0.12             | 0.1              | 116.85                   | 0.07             |
| AIFCE 9        | 53.51                   | 0.18             | 0.14             | 165.29                   | 0.11             |
| AIFCE 10       | 63.8                    | 0.24             | 0.17             | 199.05                   | 0.13             |
| AIFCE 11       | 64.56                   | 0.23             | 0.17             | 200.80                   | 0.13             |
| AIFCE 12       | 80.64                   | 0.3              | 0.22             | 249.11                   | 0.16             |
| AIFCE 13       | 77.06                   | 0.28             | 0.21             | 239.33                   | 0.15             |
| AIFCE 14       | 76.93                   | 0.29             | 0.21             | 238.66                   | 0.15             |
| AIFCE 15       | 61.91                   | 0.21             | 0.17             | 188.40                   | 0.12             |
| AIFCE 16       | 54.41                   | 0.2              | 0.15             | 170.86                   | 0.11             |
| AIFCE 17       | 31.51                   | 0.13             | 0.09             | 104.08                   | 0.06             |
| AIFCE 18       | 75.43                   | 0.28             | 0.2              | 233.38                   | 0.15             |
| AIFCE 19       | 64.18                   | 0.23             | 0.17             | 200.82                   | 0.13             |
| AIFCE 20       | 58.3                    | 0.2              | 0.16             | 180.57                   | 0.12             |
| AIFCE 21       | 64.33                   | 0.24             | 0.17             | 202.76                   | 0.13             |
| AIFCE 22       | 57.1                    | 0.22             | 0.15             | 173.62                   | 0.11             |
| AIFCE 23       | 63.2                    | 0.22             | 0.17             | 194.85                   | 0.13             |
| AIFCE 24       | 69.26                   | 0.26             | 0.19             | 215.89                   | 0.14             |
| AIFCE 25       | 65.08                   | 0.22             | 0.18             | 199.71                   | 0.13             |
| AIFCE 26       | 81.01                   | 0.31             | 0.22             | 246.88                   | 0.16             |
| AIFCE 27       | 76.39                   | 0.27             | 0.21             | 235.24                   | 0.15             |
| AIFCE 28       | 51.49                   | 0.2              | 0.14             | 163.36                   | 0.10             |
| AIFCE 29       | 15.03                   | 0.05             | 0.04             | 50.38                    | 0.03             |
| AIFCE 30       | 64.98                   | 0.23             | 0.18             | 203.99                   | 0.13             |
| <b>Minimum</b> | <b>15.03</b>            | <b>0.05</b>      | <b>0.04</b>      | <b>50.38</b>             | <b>0.03</b>      |
| <b>Maximum</b> | <b>84.34</b>            | <b>0.31</b>      | <b>0.23</b>      | <b>256.55</b>            | <b>0.17</b>      |
| <b>Mean±SD</b> | <b>59.79±16.98</b>      | <b>0.22±0.06</b> | <b>0.16±0.05</b> | <b>186.00±50.65</b>      | <b>0.12±0.03</b> |

**Table 3.** Activity concentration of naturally occurring radionuclides, absorbed dose and annual effective dose in soil samples from Federal Polytechnic Nekede.

| Sample Code      | <sup>40</sup> K (Bqkg <sup>-1</sup> ) | <sup>226</sup> Ra (Bqkg <sup>-1</sup> ) | <sup>232</sup> Th (Bqkg <sup>-1</sup> ) | Absorbed Dose(nGy <sup>-1</sup> ) | Effective Dose (μSv <sup>-1</sup> ) |
|------------------|---------------------------------------|---|---|-----------------------------------|-------------------------------------|
| FPN 1            | 101.13±1.27                           | 10.13±2.18                              | 29.11±0.71                              | 27.94                             | 34.3                                |
| FPN 2            | 86.18±1.53                            | 24.01±3.70                              | BDL                                     | 16.89                             | 20.73                               |
| FPN 3            | 70.23±1.24                            | 16.51±2.50                              | 19.70±0.71                              | 23.11                             | 28.36                               |
| FPN 4            | 124.11±1.29                           | 19.45±3.11                              | 37.15±0.81                              | 38.24                             | 46.92                               |
| FPN 5            | 90.33±1.67                            | 21.86±4.41                              | 24.90±0.83                              | 29.7                              | 36.45                               |
| FPN 6            | 99.51±1.72                            | 28.73±4.10                              | BDL                                     | 19.48                             | 23.91                               |
| FPN 7            | 72.39±1.54                            | 31.11±4.28                              | 29.05±0.59                              | 35.63                             | 43.72                               |
| FPN 8            | 100.75±1.30                           | 23.48±3.04                              | 25.91±0.73                              | 31.51                             | 38.67                               |
| FPN 9            | 98.37±1.45                            | 16.90±2.84                              | 28.19±0.84                              | 30.11                             | 36.95                               |
| FPN 10           | 87.43±1.61                            | 25.73±3.55                              | 15.48±0.57                              | 24.99                             | 30.67                               |
| FPN 11           | 93.78±1.68                            | BDL                                     | 19.11±0.68                              | 18.24                             | 22.39                               |
| FPN 12           | 72.71±1.28                            | 12.54±2.57                              | 25.86±0.77                              | 25.6                              | 31.42                               |
| FPN 13           | 110.50±1.34                           | 16.97±3.50                              | 13.99±0.53                              | 21.26                             | 26.09                               |
| FPN 14           | 113.96±1.47                           | 20.78±3.94                              | 20.17±0.82                              | 27.13                             | 33.29                               |
| FPN 15           | 147.18±1.51                           | 18.97±2.92                              | 18.57±0.59                              | 26.72                             | 32.79                               |
| FPN 16           | 89.98±1.62                            | 14.12±2.10                              | 27.51±0.71                              | 28.11                             | 34.5                                |
| FPN 17           | 103.74±1.23                           | 25.10±3.43                              | 7.80±0.53                               | 20.34                             | 24.96                               |
| FPN 18           | 59.58±1.69                            | BDL                                     | 15.67±0.66                              | 14.49                             | 17.79                               |
| FPN 19           | 63.55±1.65                            | 19.93±2.48                              | 25.10±0.74                              | 27.86                             | 34.19                               |
| FPN 20           | 110.04±1.31                           | 20.15±3.12                              | 27.81±0.71                              | 31.75                             | 38.96                               |
| FPN 21           | 98.49±1.47                            | 9.47±2.76                               | 19.88±0.83                              | 21.44                             | 26.31                               |
| FPN 22           | 114.40±1.61                           | 21.47±3.84                              | 5.44±0.54                               | 17.69                             | 21.71                               |
| FPN 23           | 69.70±1.66                            | 18.74±2.58                              | 18.37±0.68                              | 23.16                             | 28.42                               |
| FPN 24           | 105.20±1.44                           | 17.70±3.01                              | 29.86±0.71                              | 31.85                             | 39.09                               |
| FPN 25           | 98.94±1.52                            | 14.33±3.27                              | 15.92±0.64                              | 20.91                             | 25.66                               |
| FPN 26           | 104.56±1.61                           | 29.50±4.11                              | 27.70±0.57                              | 35.43                             | 43.48                               |
| FPN 27           | 78.75±1.70                            | 23.10±2.76                              | 20.18±0.80                              | 26.61                             | 32.66                               |
| FPN 28           | 88.23±1.64                            | 30.11±3.91                              | 16.99±0.75                              | 27.9                              | 34.24                               |
| FPN 29           | 104.40±1.59                           | 27.12±2.45                              | 29.17±0.66                              | 35.38                             | 43.42                               |
| FPN 30           | 30.90±1.47                            | 15.39±3.29                              | 31.49±0.83                              | 28.75                             | 35.28                               |
| <b>Minimum</b>   | <b>30.90±1.47</b>                     | <b>BDL</b>                              | <b>BDL</b>                              | <b>14.49</b>                      | <b>17.79</b>                        |
| <b>Maximum</b>   | <b>147.18±1.51</b>                    | <b>31.11±4.28</b>                       | <b>37.15±0.81</b>                       | <b>38.24</b>                      | <b>46.92</b>                        |
| <b>Mean ± SD</b> | <b>92.97±1.50</b>                     | <b>20.48±3.21</b>                       | <b>22.36±0.70</b>                       | <b>26.27±5.99</b>                 | <b>32.24±7.35</b>                   |

**Table 4.** Radium equivalent, hazard indices, annual gonadal dose equivalent and excess lifetime cancer risk in soil samples from Federal Polytechnic Nekede.

| Sample ID      | Ra (Bqkg <sup>-1</sup> ) | H <sub>in</sub>  | H <sub>ex</sub>  | AGDE (μSv <sup>-1</sup> ) | ELCR             |
|----------------|--------------------------|------------------|------------------|---------------------------|------------------|
| FPN 1          | 59.54                    | 0.19             | 0.16             | 184.74                    | 0.12             |
| FPN 2          | 36.98                    | 0.16             | 0.10             | 119.77                    | 0.07             |
| FPN 3          | 50.09                    | 0.18             | 0.14             | 155.41                    | 0.10             |
| FPN 4          | 82.13                    | 0.27             | 0.22             | 254.36                    | 0.16             |
| FPN 5          | 64.42                    | 0.23             | 0.17             | 199.99                    | 0.13             |
| FPN 6          | 42.73                    | 0.19             | 0.12             | 138.54                    | 0.08             |
| FPN 7          | 78.23                    | 0.30             | 0.21             | 240.29                    | 0.15             |
| FPN 8          | 68.29                    | 0.25             | 0.18             | 212.49                    | 0.14             |
| FPN 9          | 64.79                    | 0.22             | 0.17             | 200.94                    | 0.13             |
| FPN 10         | 54.60                    | 0.22             | 0.15             | 171.67                    | 0.11             |
| FPN 11         | 38.20                    | 0.11             | 0.10             | 120.61                    | 0.08             |
| FPN 12         | 55.12                    | 0.18             | 0.15             | 169.67                    | 0.11             |
| FPN 13         | 45.48                    | 0.17             | 0.12             | 145.61                    | 0.09             |
| FPN 14         | 58.40                    | 0.21             | 0.16             | 184.30                    | 0.12             |
| FPN 15         | 56.86                    | 0.20             | 0.15             | 182.45                    | 0.11             |
| FPN 16         | 60.39                    | 0.20             | 0.16             | 186.88                    | 0.12             |
| FPN 17         | 44.24                    | 0.19             | 0.12             | 142.74                    | 0.09             |
| FPN 18         | 30.65                    | 0.09             | 0.08             | 95.49                     | 0.06             |
| FPN 19         | 60.72                    | 0.22             | 0.16             | 186.46                    | 0.12             |
| FPN 20         | 68.39                    | 0.24             | 0.18             | 213.06                    | 0.14             |
| FPN 21         | 45.48                    | 0.15             | 0.12             | 143.29                    | 0.09             |
| FPN 22         | 38.06                    | 0.16             | 0.10             | 125.00                    | 0.08             |
| FPN 23         | 50.38                    | 0.19             | 0.14             | 156.58                    | 0.10             |
| FPN 24         | 68.50                    | 0.23             | 0.18             | 212.54                    | 0.14             |
| FPN 25         | 44.71                    | 0.16             | 0.12             | 141.89                    | 0.09             |
| FPN 26         | 77.16                    | 0.29             | 0.21             | 239.77                    | 0.15             |
| FPN 27         | 58.02                    | 0.22             | 0.16             | 180.46                    | 0.11             |
| FPN 28         | 61.20                    | 0.25             | 0.17             | 191.76                    | 0.12             |
| FPN 29         | 76.87                    | 0.28             | 0.21             | 238.51                    | 0.15             |
| FPN 30         | 62.80                    | 0.21             | 0.17             | 188.89                    | 0.12             |
| <b>Minimum</b> | <b>30.65</b>             | <b>0.09</b>      | <b>0.08</b>      | <b>95.49</b>              | <b>0.06</b>      |
| <b>Maximum</b> | <b>82.13</b>             | <b>0.30</b>      | <b>0.22</b>      | <b>254.36</b>             | <b>0.16</b>      |
| <b>Mean±SD</b> | <b>56.78±13.12</b>       | <b>0.21±0.05</b> | <b>0.15±0.04</b> | <b>177.47±39.20</b>       | <b>0.11±0.03</b> |

## DISCUSSION

The activity concentrations of primordial radionuclides have been carried out in this study. The mean activity concentrations of radionuclides in soil samples are  $88.41\pm1.51$  Bqkg<sup>-1</sup>,  $20.69\pm3.56$  Bqkg<sup>-1</sup>,  $25.04\pm0.71$  Bqkg<sup>-1</sup> for <sup>40</sup>K, <sup>226</sup>Ra, and <sup>232</sup>Th respectively, as obtained from the AIFCE and  $92.97\pm1.50$  Bqkg<sup>-1</sup>,  $20.48\pm3.21$  Bqkg<sup>-1</sup>,  $22.36\pm0.70$  Bqkg<sup>-1</sup> for <sup>40</sup>K, <sup>226</sup>Ra and <sup>232</sup>Th, respectively as obtained from FPN. These values are comparable to what was obtained from similar tertiary institutions within Nigeria, as shown in table 5. For instance, the research conducted by Eke *et al.* (8) on soil samples at the Federal University of Technology, Owerri, revealed the mean activity concentrations of  $90.18$  Bqkg<sup>-1</sup>,  $17.88$  Bqkg<sup>-1</sup> and  $22.82$  Bqkg<sup>-1</sup> for <sup>40</sup>K, <sup>226</sup>Ra and <sup>232</sup>Th, respectively. Similarly, the study conducted by Egunnyinka *et al.* (20) on evaluating primordial radionuclides in the topsoil of the University of Ibadan showed the activity concentrations  $261.37\pm192.17$ ,  $50.01\pm29.00$  and

$84.66\pm37.88$  Bqkg<sup>-1</sup> for <sup>40</sup>K, <sup>226</sup>Ra(<sup>238</sup>U) and <sup>232</sup>Th respectively. Their results are comparable to the findings in the present study.

The activity concentrations of the present study are compared with similar studies in other countries, as shown in table 5. The average activity concentrations of <sup>40</sup>K and <sup>226</sup>Ra in AIFCE and FPN are much lower than in East China, South India, Egypt, Tanzania and Iraq. However, the activity concentrations of <sup>40</sup>K from Tanzania and East China are higher than the average world values of 420, 32, and 45 Bqkg<sup>-1</sup> for <sup>40</sup>K, <sup>226</sup>Ra and <sup>232</sup>Th, respectively, as given in the UNSCEAR report (1) (table 5). The activity concentrations of <sup>226</sup>Ra in the listed countries are higher than the average world value except for Egypt and East China. The activity concentrations of <sup>232</sup>Th obtained from the present study and East China, Egypt, Tanzania, and Iraq are lower than the average world value as stated in the UNSCEAR report (1).

The results as presented in tables 1 and 3 indicate that the distribution of activity concentrations of primordial radionuclides in the studied areas are not uniform. The non-uniformity may be partly due to the geochemical, chemical, mineralogical and physical properties of the terrestrial soil and other infrastructural projects that have been ongoing since the establishment of the institutions. The migration of weathered materials from surrounding rocks might have contributed to the enhanced activity concentrations recorded in the studied areas (21, 22).

The present investigation revealed that <sup>40</sup>K contributes a significant amount to the total radioactivity of soil in AIFCE and FPN. The high value of <sup>40</sup>K could be due to potash feldspar minerals present in soil samples of the studied locations (8, 23). Moreover, the activity concentrations of <sup>40</sup>K from the present study is found to be higher than those of <sup>226</sup>Ra and <sup>232</sup>Th, which corroborates with the findings of similar studies within Nigeria.

The average absorbed dose estimated for soil samples from the present study due to primordial radionuclides was  $28.75\pm10.39$  nGyh<sup>-1</sup> and  $26.27\pm5.99$  nGyh<sup>-1</sup> AIFCE and FPN, respectively. These values are below the average world value of 59 nGyh<sup>-1</sup> for an outdoor absorbed dose as reported in UNSCEAR (1) report, indicating that the soil samples within the studied areas are free of radiological threat and they are within the normal background environment. The comparison of the absorbed dose obtained from the two institutions with existing literature indicates that the present values are less than the average value of  $32.17$  nGyh<sup>-1</sup> in Niger Delta, Nigeria (24),  $38.7\pm5.0$  nGyh<sup>-1</sup> in Agbara, Ogun State, Nigeria (25) and  $42.94$  nGyh<sup>-1</sup> in the non-oil region of Yemen (26).

Other radiological parameters such as radium equivalent activity, external and internal hazard indices have their average values  $59.79\pm16.98$  Bqkg<sup>-1</sup>,  $0.16\pm0.05$ ,  $0.22\pm0.06$ , respectively, for the AIFCE,

which are below the recommended safe limit based on UNSCEAR (1) report. Similarly, radium equivalent activity and external and internal hazard indices obtained from FPN have average values of  $56.78 \pm 13.12 \text{ Bq kg}^{-1}$ ,  $0.15 \pm 0.04$ ,  $0.21 \pm 0.05$ , respectively.

Evaluation of excess lifetime cancer risk (ELCR) revealed that radiation levels from  $^{40}\text{K}$ ,  $^{226}\text{Ra}$ , and  $^{232}\text{Th}$  in the studied soil are unlikely to cause cancer. The calculated ELCR ranged from 0.03 to 0.17 with a mean of  $0.12 \pm 0.03$  for soil samples from AIFCE, while ELCR for soil samples in FPN ranged from 0.06 to 0.16 with a mean of  $0.11 \pm 0.03$ . The estimated ELCR from the two locations is less than the average world value of  $0.290 \times 10^{-3}$  as reported by (1,3), indicating the probability of developing radiation-induced cancer from exposure from soil samples over a lifetime exposure of 70 years is low.

## CONCLUSION

The gamma-ray spectrometry method has been used to determine the activity concentrations and estimate the radiological health risks associated with soil samples from the AIFCE and FPN, Owerri. The results showed that the activity concentrations in soil samples from the campuses are comparable to the reports from other higher institutions in Nigeria. Moreover, the activity concentrations are below the average world value, as recommended by UNSCEAR (1). The activity concentrations obtained were used to calculate the absorbed dose rate, annual effective dose equivalent, annual gonadal dose and radium equivalent. The results are below the permissible safe limit.

Furthermore, excess lifetime cancer risks showed that the probability of developing radiation-induced cancer due to exposure to gamma radiation from the naturally occurring radioactive materials in soil samples is insignificant. This suggests that soil samples from the Alvan Ikoku Federal College of Education and Federal Polytechnic Nekede, Owerri are free from radioactive contamination and do not pose a threat to the two communities. However, this study's result represents reference information on radiation dose levels. It could serve as baseline data on the natural radioactivity level and epidemiological studies of the AIFCE and FPN.

## ACKNOWLEDGEMENTS

The authors wish to thank the Coordinator, Radiation and Health Physics Research Laboratory, Department of Physics, University of Ibadan, Ibadan, Nigeria, for the permission to access the research laboratory and use the thallium sodium iodide [NaI(Tl)] detector for the analysis of the soil samples.

**Conflicts of Interests:** The authors wish to state that

there are no known conflicts of interest associated with this publication.

**Funding:** The authors received no financial support for the work.

**Authors' contribution:** B.C.E designed, conceptualized the research and edited the manuscript U.M.U supervised the work I.R.A analyzed the data, wrote the manuscript and edited the manuscript All authors read and approved the manuscript.

## REFERENCES

1. UNSCEAR (2000) United Nations Scientific Committee on the effect of Atomic Radiation: Exposures from natural radiation sources. Report to General Assembly, with Scientific Annexes. United Nations, New York
2. Ramasamy V, Suresh G, Meenakshisundaram V, Gajendran V (2009) Evaluation of natural radionuclide content in river sediments and excess lifetime cancer risk due to gamma radioactivity. *Res J Environ Earth Sci*, **1**: 6-10.
3. Taskin, H, Karavus M, Ay P, Topuzoglu A, Hindiroglu S, Karahan G (2009) Radionuclide concentrations in soil and lifetime cancer risk due to the gamma radioactivity in Kirkclareli, Turkey. *J Environmental Radioactivity*, **100**: 49-53.
4. Emelue HU, Jibiri NN, Eke BC (2014) Excess Lifetime Cancer Risk due to Gamma Radiation in and Around Warri Refining and Petrochemical Company in Niger Delta, Nigeria. *British Journal of Medicine & Medical Research*, **4**(13): 2590-2598.
5. Brenner JD and Sachs RK (2006) Estimating radiation-induced cancer risk at very low doses: rationale for using a linear no-threshold approach. *Radiation Environ Biology*, **44**: 253-256.
6. Ibrahim FA and Mohammad IA (2009) Soil radioactivity levels and radiation hazard assessment in the highlands of Northern Jordan. *Radiation Measurements*, **44**: 102-110.
7. Mehta R, Badhan K, Sonkawade RG, kansal S, Singh S (2010) Analysis of terrestrial natural radionuclides in soil samples and assessment of average effective dose. *Indian Journal of Pure and Applied Physics*, **48**: 805-808.
8. Eke BC, Jibiri NN, Anusionwu BC, Orji CE, Emelue HU (2015) Baseline Measurements of natural radioactivity in soil samples from the Federal University of Technology, Owerri, South-East, Nigeria. *British J Applied Sciences & Technology*, **5**(2): 142 -149.
9. Obed RI, Farai IP, Jibiri NN (2005) Population dose distribution due to soil radioactivity concentration levels in 18 cities across. *Nigeria J Radiat Prot*, **25**: 305-312.
10. Jibiri NN and Akomolafe IR (2016). Radiological assessment and geochemical characterization of the sediments of Awba Dam, University of Ibadan, Nigeria. *Radiat Prot Environ*, **39**: 222-32.
11. Jibiri NN and Emelue HU (2008) Soil radionuclide concentrations and radiological assessment in and around a refining and petrochemical company in Warri, Niger Delta, Nigeria. *Journal of Radio logical Protection*, **28**: 361-368.
12. Beretka J and Matthew PJ (1985) Natural radioactivity of Australian building materials, industrial waste, and by-products. *Health Physics*, **48**: 87-95.
13. Farai IP, Obed RI, Jibiri NN (2006) Soil radioactivity and incidence of cancer in Nigeria. *Journal of Environmental Radioactivity*, **90**: 29 -36.
14. Jibiri NN and Okeyode I (2011) Activity concentrations of natural radionuclides in the sediments of Ogun River, Southwestern Nigeria. *Radiation Protection Dosimetry*, **147**(4): 555-64.
15. Nwankwo CU, Ogundare FO, Folley DE (2015) Radioactivity concentration variation with depth and assessment of worker's doses in selected mining sites. *J Radiat Res Appl Sci*, **8**: 216-220.
16. UNSCEAR (1988) United Nations Scientific Committee on the Effect of Atomic Radiation Sources, effects and risks of ionizing radiation report to the general assembly with annexes. United States publication E88ix 17 (United Nations) New York.
17. Ridha AA and Hasan HA (2016) Cancer Risk Due to the Natural Radioactivity in Cigarette Tobacco. *Detection*, **4**: 54-65. <http://dx.doi.org/10.4236/detection.2016.43008>.
18. International Commission on Radiological Protection, ICRP (1991). *Series Report of the 1990 recommendations of the ICRP, Publication No. 60. Ann. ICRP*, **21**: 1-3.

19. Kolo MT, Amin YM, Khandaker MU, Abdullah WHB (2017) Radio-nuclide concentrations and excess lifetime cancer risk due to gamma radioactivity in tailing enriched soil around Maiganga coalmine, Northeast Nigeria. *Int J Radiat Res*, **15** (1): 71-80.

20. Egunyinka OA, Olowookere CJ, Jibiri NN, Babalola IA, Obed RI (2009) An evaluation of  $^{238}\text{U}$ ,  $^{40}\text{K}$ , and  $^{232}\text{Th}$  concentrations in the top soil of the university of Ibadan (UI), Southwestern Nigeria. *The Pacific Journal of Science and Technology*, **10**: 742-752.

21. Suresh GM, Ravisankar R, Rajalakshmi A, Sivakumar S, Chandrasekaran A, Anand DP (2014) Measurements of natural gamma radiation in beach sediments of north east coast of Tamilnadu, India by gamma-ray spectrometry with multivariate statistical approach. *J Radiat Res Appl Sci*, **7**(1): 7-17.

22. Ugbede FO, Okoye ONN, Akpolile AF, Oladele BB (2021) Baseline Radioactivity in the Soil of Evangel Take-Off Campus, Evangel University, Nigeria, and its Associated Health Risks. *Chemistry Africa*, **4**: 703-713.

23. Arafa W (2004) Specific activity and hazards of granite samples collected from the Eastern Desert of Egypt. *J Environ Radioact*, **75** (3): 315-327.

24. Avwiri GO and Ononugbo CP (2012) Natural radioactivity levels in surface soil of Ogba/Egbema/Ndoni oil and gas fields. *Energy Sci Technol*, **4**(2): 92-101.

25. Gbadamosi MR, Banjoko OO, Abudu KA, Ogunbanjo OO, Ogunneye AL (2017) Radiometric evaluation of excessive lifetime cancer probability due to naturally occurring radionuclides in wastes dumpsites soils in Agbara, Southwest, Nigeria. *J Assoc Arab Uni Basic Appl Sci*, **24**: 315-324.

26. Ali MMM, Zhao H, Rawashdeh A, Mohammed YA, Al Hassan M (2021) Assessment of radiation hazard indices for sand samples from Ma'rib in Yemen. *Int J Radiat Res*, **19**(3): 615-623.

27. Aladeniyi K, Olowookere C, Oladele BB (2019) Measurement of natural radioactivity and radiological hazard evaluation in the soil samples collected from Owo, Ondo State, Nigeria. *J Radiat Res Appl Sci*, **12**(1): 200-209.

28. Olagbaju PO, Okeyode IC, Alatise OO, Bada BS (2021) Background radiation level measurement using hand-held dosimeter and gamma spectrometry in Ijebu-Ife, Ogun State Nigeria. *Int J Radiat Res*, **19**(3): 591-598.

29. Akpan AE, Paul ND, Uwah EJ (2016) Ground radiometric investigation of natural radiation levels and their radiological effects in Akpabuyo, Nigeria. *J Afr Earth Sci*, **123**: 185-192.

30. Zhou Z, Yang Z, Sun Z, Liao Q, Guo Y, Chen J (2020) Multidimensional pollution and potential ecological and health risk assessments of radionuclides and metals in the surface soils of a uranium mine in East China. *J Soils Sediments*, **20**: 775-791.

31. Sivakumar R (2014) An assessment of natural radioactivity levels and radiation hazards in the soil of Coonoor, South India. *Environ Earth Sci*, **72**: 5063-5071.

32. Ebaid YY, El-Tahawy MS, El-Lakany AA, Garcia SR, Brooks GH (2000) Environmental radioactivity measurements of Egyptian soils. *J Radioanal Nucl Chem*, **243**(2): 543-550.

33. Mohammed NK and Mazunga MS (2013) Natural radioactivity in soil and water from Likuya village in the neighborhood of Mkuju uranium deposit. *Int J Analyt Chem*, **2013**: 501856.

34. Taqi AH, Shaker AM, Battawy AA (2018) Natural radioactivity assessment in soil samples from Kirkuk city of Iraq using HPGe detector. *Int J Radiat Res*, **16**(4): 455-463.