

# Potential radiological risks associated with artisanal mining activities and the use of tailings from Kuru District as building materials

M.B. Adedokun<sup>1\*</sup>, A.Z. Ibitoye<sup>2</sup>, K.I. Ogungbemi<sup>1</sup>

<sup>1</sup>Department of Physics, Faculty of Science, University of Lagos, Akoka, Lagos, Nigeria

<sup>2</sup>Department of Radiation Biology, Radiotherapy, and Radiodiagnosis, College of Medicine, University of Lagos, Idi-Araba, Lagos, Nigeria

## ABSTRACT

### ► Original article

#### \*Corresponding author:

Margaret Bose Adedokun, Ph.D.,

#### E-mail:

madedokun@unilag.edu.ng

Received: March 2024

Final revised: October 2024

Accepted: November 2024

Int. J. Radiat. Res., July 2024;  
23(3): 593-600

DOI: 10.61186/ijrr.23.3.12

**Keywords:** Mining activities, tailings, building material, radiation hazard indices, gamma spectrometry.

**Background:** This study evaluated the radiological risks associated with mining and living in buildings constructed with tailings. **Materials and Methods:** A NaI detector was used to determine the specific activity of <sup>238</sup>U, <sup>232</sup>Th, and <sup>40</sup>K in tailings, and radiological hazard indices were estimated. **Results:** The mean specific activities of <sup>238</sup>U, <sup>232</sup>Th, and <sup>40</sup>K were 84.27 ± 13.62 Bq/kg, 11.65 ± 0.69 Bq/kg, and 193.26 ± 9.98 Bq/kg, respectively. The mean values for <sup>232</sup>Th and <sup>40</sup>K were about 74% and 54% lower than their corresponding average world values, while that of <sup>238</sup>U was 155% higher than its average world value. The hazards indices: radium equivalent (Ra<sub>eq</sub>), absorbed dose rate (D<sub>out</sub> and D<sub>in</sub>), annual effective dose equivalent (ADE<sub>out</sub> and ADE<sub>in</sub>), annual gonadal dose equivalent (AGDE), external hazard index (H<sub>ex</sub>), internal hazard index (H<sub>in</sub>), and excess lifetime cancer risk (ELCR<sub>out</sub> and ELCR<sub>in</sub>) were estimated from the specific activity of the radionuclides. Their mean values were 249.74 Bq/kg, 54.03 nGy/h, 106.00 nGy/h, 66.31 μSv/y, 520.35 μSv/y, 369.78 μSv/y, 0.31, 0.54, 0.232 × 10<sup>-3</sup> and 1.821 × 10<sup>-3</sup> respectively. Ra<sub>eq</sub>, D<sub>out</sub>, ADE<sub>out</sub>, H<sub>ex</sub> and H<sub>in</sub> were about 32%, 5%, 17.11%, 69%, and 46% lower, while D<sub>in</sub>, ADE<sub>in</sub>, and AGDE were about 26.19%, 24%, and 24.09% higher than their respective reference values. ELCR<sub>out</sub> is about 80%, while ELCR<sub>in</sub> is almost six times higher than the reference ELCR from all carcinogens. **Conclusion:** The high specific activity of <sup>238</sup>U, D<sub>in</sub>, ADE<sub>in</sub>, AGDE, and ELCR<sub>in</sub> indicates that tailings are unsuitable for building and that there are potential radiological risks associated with artisanal mining activities in Kuru district.

## INTRODUCTION

Ionizing radiation is an intrinsic component of the Earth's environment, and humans have been exposed to it throughout history <sup>(1)</sup>. Environmental radioactivity arises from natural radioactive materials present on Earth, either through natural processes or anthropogenic activities. Radiation levels can be influenced by geological formation, soil mineral concentrations <sup>(2)</sup>, and human activities such as nuclear power, nuclear research, radiography, cement and paint production, fertiliser manufacturing, oil exploration, and mining <sup>(3)</sup>. Workers in these industries are subject to continuous radiation exposure, primarily through inhalation and dermal contact.

In addition, residential, and occupational environments, such as houses and offices, can contribute significantly to external radiation exposure. This is largely due to gamma-ray emissions from building materials that contain radionuclides such as <sup>238</sup>U, <sup>232</sup>Th, and <sup>40</sup>K and their decay products <sup>(4)</sup>. While the importance of shelter cannot be overemphasised, building constructed with materials

containing high levels of radiation will contribute significantly to the radiation exposure of the occupants. Various construction materials contain differing levels of naturally occurring radionuclides that release radon <sup>(5)</sup>.

Continuous exposure to radon, a decay product of <sup>238</sup>U, affects indoor air quality and has been linked to the induction of lung cancer <sup>(6)</sup>. Therefore, building materials with higher levels of naturally occurring radionuclides will lead to increased radon concentrations over time as these radionuclides accumulate. It is essential to assess the levels of radionuclides in building materials to evaluate the population exposure, especially given that people spend approximately 78% of their time indoors <sup>(7)</sup>.

Mining activities in Jos began around 1904, and the city has long been known for its mining activities. Although formal mining operations have ceased, illegal artisanal mining continues <sup>(8)</sup>. This type of mining is associated with serious risks, including fatalities, injuries, and significant environmental disruptions <sup>(9-12)</sup>. Despite these dangers, artisanal mining remains a vital means of livelihood in the region <sup>(13)</sup>, with its economic benefits often

outweighing concerns about its consequences.

In the Kuru District, mineral ore extraction is now largely done through manual centrifugation, with mining tailings (waste products) left in piles around the mining sites. These tailings contain various minerals, some of which are carcinogenic (14). These soil-rich wastes materials are often collected for use in building local ovens and shelters, as they are inexpensive (14). While the radiological effects of mining activities in Jos have been widely studied-focusing on population exposure through water consumption, the food chain (11,15), dose rates in the air at active mining sites (16), and the impact of tin mining on different soil types and concrete blocks (17, 18), there is limited research on the radiological risks associated with artisanal mining. Specifically, little attention has been given to the risks faced by residents living in homes built with mining tailings. This study aims to assess the potential long-term radiological risks posed by artisanal mining activities in the Kuru District, focusing on miners and residents living in homes constructed using tailings from mining sites in Jos.

## MATERIALS AND METHODS

### Study area

This study was conducted in three mining locations: Kuru-Jenta (KJ), Kanikon (KK) and Zakong-Soil (ZS), all located in the Kuru District (figure 1) of Jos South Local Government Area of Plateau State. The area is situated at an elevation of about 1318.20 m, with coordinates at 9.7184° North and 8.8359° East.

While agriculture is the main occupation in Jos, Plateau State is widely known for its mining industry. Historically, it has been Nigeria's leading region for mining, especially in the extraction and exportation of tin and columbite. In the Kuru District, miners work manually digging pits with basic tools like shovels, diggers, and spades to extract minerals. The minerals mined here include cassiterite, tin, wolframite, ilmenite, and columbite.




Figure 1. Map of Kuru District and its environs (adapted from (19)).

### Collection and preparation of samples

A total of 21 samples were collected from the three mining locations. Six samples were obtained

from Kuru-Jenta (KJ), five from Kanikon (KK), and ten from Zakong-soil (ZS). The number of samples collected at each site was proportional to the area's size. Approximately 1 kg of tailings was collected from the surface (about 0-30 cm) of randomly selected heaps at each location. The samples were placed in plastic bags, clearly labelled, and transported to the laboratory. Stones were manually removed from each sample before placing them in pre-cleaned plastic trays. The samples were air-dried in the laboratory for six weeks to ensure all moisture was removed. After drying, the samples were pulverised using a laboratory mortar and pestle and then passed through a 1 mm sieve for homogeneity. About 300 g of each sieved sample was transferred to a clean, pre-weighed cylindrical plastic container. The filled containers were weighed to determine the exact mass of each sample. All containers were hermetically sealed and left for four weeks to allow the  $^{238}\text{U}$  and  $^{232}\text{Th}$  and their progenies to reach secular equilibrium.

### Sample measurement and analysis of spectral.

The specific activities of  $^{238}\text{U}$ ,  $^{232}\text{Th}$ , and  $^{40}\text{K}$  in the tailing samples were measured using a gamma-ray spectrometer. This spectrometer was equipped with a 76 mm by 76 mm sodium iodide doped with thallium (NaI (Tl)), Model 802-series (Canberra inc., USA). The NaI (Tl) detector was shielded with thick lead blocks and connected to a Canberra Multichannel Analyser (MCA), Model 2007P, through a pre-amplifier base for data acquisition. The operating voltage of the detector was set to 600 V with a resolution of 7.5% full width half maximum (FWHM) at the 0.662 MeV peak of  $^{137}\text{Cs}$ . Energy and efficiency calibrations were performed using gamma sources provided by the International Atomic Energy Agency (IAEA), following IAEA Document No. 385 (20).

To determine the laboratory's background radiation, an empty beaker identical in size to the sample beaker was placed in the detector and counted under the same conditions. Each sample was then placed in the detector and counted for 25200 s (7 h). The net counts for each sample were obtained by subtracting the background counts, and the spectra were analysed using GENIE-2000 software (Canberra Inc., USA). The specific activities of  $^{238}\text{U}$ ,  $^{232}\text{Th}$ , and  $^{40}\text{K}$  were calculated based on the concentrations of  $^{214}\text{Bi}$  (at the 1764.5 keV gamma peak),  $^{208}\text{Tl}$  (2614.7 keV gamma peak), and the gamma peak of  $^{40}\text{K}$  (1460.8 keV), respectively. The minimum detectable activity (MDA) of the detector for the chosen gamma peaks were calculated from the background radiation count rate, as as described in literature (21, 22), and is presented in table 1.

### Specific activity (A)

The specific activity A (Bq/kg) for all samples was determined using equation 1 (20, 23).

$$A = \frac{C}{t \times m \times \epsilon \times P_Y} \quad (1)$$

Where; A is the specific activity of the radionuclide in Bq/kg, C is the net count for the sample in the peak energy range, m is the mass of the sample in kg, t is the counting lifetime in seconds,  $\epsilon$  is the detector energy-dependent efficiency, and  $P_Y$  is the gamma-ray yield per disintegration of radionuclides.

**Table 2.** Specific activity of  $^{238}\text{U}$ ,  $^{232}\text{Th}$  and  $^{40}\text{K}$  in tailings from Kanikon (mean  $\pm$  SD).

| Sample          | $^{238}\text{U}$ (Bq/kg) | $^{232}\text{Th}$ (Bq/kg) | $^{40}\text{K}$ (Bq/kg) |
|-----------------|--------------------------|---------------------------|-------------------------|
| KK1             | 99.66 $\pm$ 16.21        | 8.81 $\pm$ 0.52           | 151.21 $\pm$ 7.97       |
| KK2             | 161.05 $\pm$ 25.59       | 6.97 $\pm$ 0.41           | 217.84 $\pm$ 11.40      |
| KK3             | 92.48 $\pm$ 14.89        | 7.16 $\pm$ 0.42           | 158.95 $\pm$ 8.28       |
| KK4             | 79.89 $\pm$ 13.13        | 11.13 $\pm$ 0.65          | 229.90 $\pm$ 11.93      |
| KK5             | 22.50 $\pm$ 4.05         | 7.16 $\pm$ 0.42           | 185.79 $\pm$ 9.74       |
| Mean            | 91.12 $\pm$ 14.77        | 8.25 $\pm$ 0.48           | 188.74 $\pm$ 9.86       |
| World's average | 33.00                    | 45.00                     | 420.00                  |
| P               | 0.058                    | 0.001                     | 0.001                   |

### Radiological hazard indices

Various radiological hazard indices were calculated to assess potential radiological risks associated with mining in Kuru and the use of tailings in building construction. These indices include radium equivalent ( $R_{\text{eq}}$ ), absorbed dose rate (D), annual effective dose equivalent (E), annual gonadal dose equivalent (AGDE), external hazard index ( $H_{\text{ex}}$ ), internal hazard index ( $H_{\text{in}}$ ), and excess lifetime cancer risk (ELCR) and have been described fully elsewhere <sup>(22)</sup>.

### Radium equivalent ( $R_{\text{eq}}$ )

Radium equivalent provides a single index to estimate the gamma radiation output from a mix of radionuclides in samples. It was calculated using equation 2 <sup>(24, 25)</sup>:

$$R_{\text{eq}} = A_{\text{U}} + 1.43A_{\text{Th}} + 0.77A_{\text{K}} \quad (2)$$

Where;  $A_{\text{U}}$ ,  $A_{\text{Th}}$ , and  $A_{\text{K}}$  are the specific activities for  $^{238}\text{U}$ ,  $^{232}\text{Th}$ , and  $^{40}\text{K}$ , respectively.

### Absorbed dose rate (D)

The outdoor absorbed dose rate in air was calculated using equation 3 <sup>(25)</sup>, while the indoor absorbed dose rate was calculated using equation 4 <sup>(26, 27)</sup>:

$$D_{\text{outdoor}}(\text{nGy h}^{-1}) = 0.462A_{\text{U}} + 0.604A_{\text{Th}} + 0.0417A_{\text{K}} \quad (3)$$

$$D_{\text{indoor}}(\text{nGy h}^{-1}) = 0.92A_{\text{U}} + 1.1A_{\text{Th}} + 0.08A_{\text{K}} \quad (4)$$

Where;  $A_{\text{U}}$ ,  $A_{\text{Th}}$ , and  $A_{\text{K}}$  are the specific activities of  $^{238}\text{U}$ ,  $^{232}\text{Th}$ , and  $^{40}\text{K}$  in Bq/kg, respectively.

### Annual effective dose (AED)

$AED_{\text{in}}$  accounts for dwellers' exposure, while  $AED_{\text{out}}$  accounts for miners' exposure. The outdoor and indoor annual effective ( $AED_{\text{out}}$  and  $AED_{\text{in}}$ ) doses were calculated using equations 5 and 6 <sup>(28)</sup>:

$$AED_{\text{out}} (\mu\text{Sv/y}) = D_{\text{out}} (\times 0.2 \times 0.7 \times 8766 \times 10^{-3}) \quad (5)$$

$$AED_{\text{in}} (\mu\text{Sv/y}) = D_{\text{in}} (\times 0.8 \times 0.7 \times 8766 \times 10^{-3}) \quad (6)$$

Where;  $D_{\text{out}}$  and  $D_{\text{in}}$  are the outdoor and indoor absorbed dose rate respectively in nGy/h, 0.2 is the outdoor occupancy, 0.8 is the indoor occupancy factor, 0.7 Sv/Gy is the conversion factor from absorbed dose rate to the effective dose rate, and 8766 is the number of hours in a year.

### Annual gonadal dose equivalent (AGDE)

The AGDE (for miners and dwellers) has been calculated using equation 7 <sup>(25, 29)</sup>:

$$AGDE (\mu\text{Sv/y}) = 3.09A_{\text{U}} + 4.18A_{\text{Th}} + 0.314A_{\text{K}} \quad (7)$$

Where;  $A_{\text{U}}$ ,  $A_{\text{Th}}$ , and  $A_{\text{K}}$  are the specific activities of  $^{238}\text{U}$ ,  $^{232}\text{Th}$ , and  $^{40}\text{K}$  in Bq/kg, respectively.

### Hazard indices ( $H_{\text{ex}}$ and $H_{\text{in}}$ )

External and internal index  $H_{\text{ex}}$  and  $H_{\text{in}}$  were calculated using the specific activities of  $^{238}\text{U}$ ,  $^{232}\text{Th}$ , and  $^{40}\text{K}$  in sampled tailings using equations 8 and 9, respectively <sup>(28)</sup>:

$$H_{\text{ex}} = \frac{A_{\text{U}}}{370 \text{ Bq/kg}} + \frac{A_{\text{Th}}}{259 \text{ Bq/kg}} + \frac{A_{\text{K}}}{4810 \text{ Bq/kg}} \quad (8)$$

$$H_{\text{in}} = \frac{A_{\text{U}}}{185 \text{ Bq/kg}} + \frac{A_{\text{Th}}}{259 \text{ Bq/kg}} + \frac{A_{\text{K}}}{4810 \text{ Bq/kg}} \quad (9)$$

Where;  $A_{\text{U}}$ ,  $A_{\text{Th}}$ , and  $A_{\text{K}}$  are the values for the specific activities of  $^{238}\text{U}$ ,  $^{232}\text{Th}$ , and  $^{40}\text{K}$ .

### Excess lifetime cancer risk (ELCR)

Excess lifetime cancer risk (ELCR) from outdoor and indoor exposures have been calculated using equations 10 and 11 <sup>(22)</sup> to estimate the potential carcinogenic effects of exposure to radionuclides by miners and dwellers resulting from mining activities and the use of tailings as a building material, respectively.

$$ELCR_{\text{out}} = AED_{\text{out}} \times \text{RF} \times \text{DL} \quad (10)$$

$$ELCR_{\text{in}} = AED_{\text{in}} \times \text{RF} \times \text{DL} \quad (11)$$

Where;  $AED_{\text{out}}$  and  $AED_{\text{in}}$  are the outdoor and indoor annual effective dose, DL is the duration of life (70 years), and RF is the fatal cancer risk factor per Sievert, which is 0.05 for the public.

### Statistical Analysis

A one-sample t-test using the Statistical Package for Social Science (SPSS) version 23 (IBM, Armonk, NY) to compare the obtained values with global reference values and other relevant data. A p-value of less than 0.05 ( $p < 0.05$ ) was considered statistically significant.

## RESULTS

The lowest level of radioactivity of  $^{238}\text{U}$ ,  $^{232}\text{Th}$ , and

<sup>40</sup>K that the detector can reliably measure under the analysis conditions are shown in Table 1. The minimum detectable activities (MDA) of <sup>238</sup>U, <sup>232</sup>Th, and <sup>40</sup>K of the detector are 0.022, 0.020, and 0.080 Bq/kg, respectively. Table 2 presents the specific activities of <sup>238</sup>U, <sup>232</sup>Th, and <sup>40</sup>K in tailings from Kanikon. The values for <sup>238</sup>U ranged from 22.50 ± 4.05 to 161.05 ± 25.59 Bq/kg, for <sup>232</sup>Th ranged from 6.97 ± 0.41 to 11.13 ± 0.65 Bq/kg, and for <sup>40</sup>K ranged from 151.21 ± 7.97 to 229.90 ± 11.93 Bq/kg. The mean values are 91.12 ± 14.77, 8.25 ± 0.48, and 188.74 ± 9.86 Bq/kg for <sup>238</sup>U, <sup>232</sup>Th, and <sup>40</sup>K in tailing from Kanikon, respectively.

For the tailings collected from Kuru-Jenta (table 3), the specific activity of <sup>238</sup>U ranged from 44.54 ± 7.03 to 366.73 ± 57.44 Bq/kg, <sup>232</sup>Th ranged from 6.68 ± 0.39 to 60.20 ± 3.46 Bq/kg, and <sup>40</sup>K ranged from 251.46 ± 13.06 to 419.87 ± 20.79 Bq/kg. The mean specific activities of <sup>238</sup>U, <sup>232</sup>Th, and <sup>40</sup>K are 125.30 ± 19.85, 18.46 ± 1.07, and 304.44 ± 15.61 Bq/kg, respectively.

Table 4 shows the specific activities of <sup>238</sup>U, <sup>232</sup>Th, and <sup>40</sup>K in tailings from Zagong-soil, where the largest number of samples were analysed. The values of <sup>238</sup>U ranged from 29.46 ± 5.05 to 90.97 ± 14.84 Bq/kg, with an average value of 55.93 ± 18.69 Bq/kg. The values of <sup>232</sup>Th ranged from 4.35 ± 0.42 to 13.82 ± 0.81 Bq/kg, and <sup>40</sup>K ranged from 4.75 ± 0.23 to 226.67 ± 11.26 Bq/kg. The mean values are 55.93 ± 18.69, 9.27 ± 0.56, and 128.82 ± 6.65 Bq/kg for <sup>238</sup>U, <sup>232</sup>Th and <sup>40</sup>K.

**Table 2.** Specific activity of <sup>238</sup>U, <sup>232</sup>Th, and <sup>40</sup>K in tailings from Kanikon (mean ± SD).

| Sample          | <sup>238</sup> U (Bq/kg) | <sup>232</sup> Th (Bq/kg) | <sup>40</sup> K (Bq/kg) |
|-----------------|--------------------------|---------------------------|-------------------------|
| KK1             | 99.66±16.21              | 8.81±0.52                 | 151.21±7.97             |
| KK2             | 161.05±25.59             | 6.97±0.41                 | 217.84±11.40            |
| KK3             | 92.48±14.89              | 7.16±0.42                 | 158.95±8.28             |
| KK4             | 79.89± 13.13             | 11.13±0.65                | 229.90±11.93            |
| KK5             | 22.50± 4.05              | 7.16±0.42                 | 185.79±9.74             |
| Mean            | 91.12±14.77              | 8.25±0.48                 | 188.74±9.86             |
| World's average | 33.00                    | 45.00                     | 420.00                  |
| P               | 0.058                    | 0.001                     | 0.001                   |

**Table 3.** Specific activity of <sup>238</sup>U, <sup>232</sup>Th, and <sup>40</sup>K in tailings from Kuru-Jenta (mean ± SD).

| Sample          | <sup>238</sup> U (Bq/kg) | <sup>232</sup> Th (Bq/kg) | <sup>40</sup> K (Bq/kg) |
|-----------------|--------------------------|---------------------------|-------------------------|
| KJ1             | 56.41±8.99               | 6.68±0.39                 | 256.47±13.23            |
| KJ2             | 97.14±15.84              | 11.42±0.67                | 251.46±13.06            |
| KJ3             | 44.54±7.03               | 9.03±0.52                 | 318.13±15.78            |
| KJ4             | 81.01±12.71              | 10.75±0.61                | 419.87±20.79            |
| KJ5             | 105.98±17.07             | 12.70±0.74                | 277.97±14.44            |
| KJ6             | 366.73±57.44             | 60.20±3.46                | 302.75±16.38            |
| Mean            | 125.30±19.85             | 18.46±1.07                | 304.44±15.61            |
| World's average | 33.00                    | 45.00                     | 420.00                  |
| P               | 0.130                    | 0.070                     | 0.019                   |


As seen in tables 2 to 4, the specific activities of <sup>238</sup>U, <sup>232</sup>Th, and <sup>40</sup>K in the samples from the study areas are compared to world averages. Figures 2, 3, and 4 highlight the distribution of <sup>238</sup>U, <sup>232</sup>Th, and <sup>40</sup>K across all sampled tailings from the three mining fields. For <sup>238</sup>U, the specific activity ranges from

22.50 ± 4.05 Bq/kg in sample KK5 to 366.73 ± 57.44 Bq/kg in sample KJ6. The specific activity of <sup>232</sup>Th varies, with the lowest being 4.35 ± 0.42 Bq/kg in sample ZS2 and the highest at 60.20 ± 3.46 Bq/kg in sample KJ6. For <sup>40</sup>K, the specific activity spans from 4.75 ± 0.23 Bq/kg in sample ZS2 to 419.87 ± 20.79 Bq/kg in sample KJ4. Figure 5 compares the overall mean specific activities of <sup>238</sup>U, <sup>232</sup>Th, and <sup>40</sup>K in all the sampled tailings from the selected mining sites in Kuru District with the world average values. The overall mean specific activities of <sup>238</sup>U, <sup>232</sup>Th, and <sup>40</sup>K are recorded as 84.27 ± 13.62, 11.65 ± 0.69, and 193.26 ± 9.98 Bq/kg, respectively.


**Table 4.** Specific activity of <sup>238</sup>U, <sup>232</sup>Th, and <sup>40</sup>K in tailings from Zagong-Soil (mean ± SD).

| Sample          | <sup>238</sup> U (Bq/kg) | <sup>232</sup> Th (Bq/kg) | <sup>40</sup> K (Bq/kg) |
|-----------------|--------------------------|---------------------------|-------------------------|
| ZS1             | 54.83±9.17               | 8.13±0.48                 | 6.26±0.32               |
| ZS2             | 32.05±5.54               | 4.35±0.42                 | 166.24±8.62             |
| ZS3             | 29.46±5.05               | 7.73±0.45                 | 87.30±4.57              |
| ZS4             | 45.58±7.22               | 8.44±0.48                 | 4.75±0.23               |
| ZS5             | 90.97±14.84              | 10.14±0.60                | 176.59±9.24             |
| ZS6             | 76.82±12.08              | 11.85±0.68                | 226.67±11.26            |
| ZS7             | 57.47±9.87               | 13.82±0.81                | 145.69±7.61             |
| ZS8             | 53.91±9.01               | 5.73±0.34                 | 121.01±6.27             |
| ZS9             | 52.97±8.95               | 9.41±0.55                 | 155.17±8.08             |
| ZS10            | 65.20±11.26              | 13.10±0.77                | 198.48±10.32            |
| Mean            | 55.93±18.69              | 9.27±0.56                 | 128.82±6.65             |
| World's average | 33.00                    | 45.00                     | 420.00                  |
| P               | 0.004                    | 0.001                     | 0.001                   |


Note: World average values refer to the global average activity concentration of radionuclides in soils as reported by UNSCEAR<sup>(28)</sup>.



**Figure 2.** Distribution of the mean specific activity of <sup>238</sup>U in all sampled tailings.



**Figure 3.** Distribution of the mean specific activity of <sup>232</sup>Th in all sampled tailings.



**Figure 4.** Distribution of the mean specific activity of <sup>40</sup>K in all sampled tailings.

Figure 5. Average specific activities of <sup>238</sup>U, <sup>232</sup>Th, and <sup>40</sup>K (Bq/kg) versus the world's average values.




Table 6 presents the values of the estimated radiological hazard indices and their corresponding reference or world's average values.  $R_{a_{eq}}$  ranges from 61.30 to 685.93 Bq/kg, with a mean of 249.74 Bq/kg. The outdoor absorbed dose rate in air  $D_{out}$  ranges from 21.92 to 218.41 nGy/h with a mean value of 54.03 nGy/h. The indoor absorbed dose rate in air  $D_{in}$  to assess the absorbed dose of residents of homes built with these tailings ranges from 42.68 to 428.13 nGy/h; the mean value is 106.00 nGy/h. The indoor annual effective dose equivalent  $AED_{in}$  has values from 209.50 to 2101.69  $\mu$ Sv/y with an average value of 520.35  $\mu$ Sv/y.

Table 5. Mean specific activities of <sup>238</sup>U, <sup>232</sup>Th, and <sup>40</sup>K (Bq/kg) in the soil of some previously studied mining sites and results from the current study.

| <sup>238</sup> U | <sup>232</sup> Th | <sup>40</sup> K | Location           | Reference     |
|------------------|-------------------|-----------------|--------------------|---------------|
| 7220.00          | 16800             | -               | Plateau, Nigeria   | [8]           |
| 55.30            | 26.40             | 505.10          | South-west Nigeria | [25]          |
| 3867.50          | 8301.90           | 1251.70         | Plateau, Nigeria   | [30]          |
| 132.60           | 351.4             | 319.60          | Plateau, Nigeria   | [31]          |
| 49.71            | 4.98              | 615.21          | Plateau, Nigeria   | [32]          |
| 84.27±<br>13.62  | 11.65±<br>0.69    | 193.26±<br>9.98 | Plateau, Nigeria   | Present study |
| p=0.204          | p=0.201           | p=0.170         |                    |               |

Table 6. Summary of some estimated radiological indices from the specific activity of <sup>238</sup>U, <sup>232</sup>Th and <sup>40</sup>K measured from sampled tailings and their corresponding reference values.

| Sample          | $R_{a_{eq}}$ (Bq/kg) | $D_{out}$ (nGy/hr) | $D_{in}$ (nGy/hr) | $AED_{out}$ ( $\mu$ Sv/y) | $AED_{in}$ ( $\mu$ Sv/y) |
|-----------------|----------------------|--------------------|-------------------|---------------------------|--------------------------|
| KK1             | 228.69               | 57.67              | 113.63            | 70.77                     | 557.79                   |
| KK2             | 338.75               | 87.70              | 173.48            | 107.62                    | 851.60                   |
| KK3             | 225.11               | 53.68              | 105.83            | 65.88                     | 519.53                   |
| KK4             | 272.83               | 53.22              | 104.36            | 65.32                     | 512.32                   |
| KK5             | 175.80               | 22.47              | 43.62             | 27.57                     | 214.15                   |
| KJ1             | 263.44               | 40.79              | 80.20             | 50.06                     | 392.81                   |
| KJ2             | 307.09               | 62.26              | 122.30            | 76.41                     | 600.36                   |
| KJ3             | 302.41               | 39.30              | 76.68             | 48.23                     | 376.41                   |
| KJ4             | 419.68               | 61.43              | 120.36            | 75.38                     | 590.86                   |
| KJ5             | 338.18               | 68.22              | 133.99            | 83.73                     | 657.74                   |
| KJ6             | 685.93               | 218.41             | 428.13            | 268.04                    | 2101.69                  |
| ZS1             | 71.28                | 30.50              | 59.89             | 37.43                     | 294.02                   |
| ZS2             | 166.28               | 24.36              | 47.74             | 29.90                     | 234.34                   |
| ZS3             | 107.73               | 21.92              | 42.68             | 26.90                     | 209.50                   |
| ZS4             | 61.30                | 26.35              | 51.60             | 32.34                     | 253.31                   |
| ZS5             | 241.44               | 55.52              | 109.15            | 68.13                     | 535.81                   |
| ZS6             | 271.30               | 53.49              | 104.83            | 65.64                     | 514.60                   |
| ZS7             | 189.41               | 40.97              | 79.88             | 50.28                     | 392.10                   |
| ZS8             | 155.28               | 33.41              | 65.70             | 41.01                     | 322.53                   |
| ZS9             | 185.91               | 36.63              | 71.65             | 44.94                     | 351.74                   |
| ZS10            | 236.76               | 46.32              | 90.47             | 56.83                     | 444.12                   |
| Mean            | 249.74               | 54.03              | 106.00            | 66.31                     | 520.35                   |
| World's average | NA                   | 57.00              | 84.00             | 80.00                     | 420.00                   |
| reference level | 370.00               | NA                 | NA                | NA                        | NA                       |
| P               | 0.001                | 0.745              | 0.228             | 0.230                     | 0.261                    |

Table 7. Measured Indoor AED (mSv/yr) from different studied building materials across the world and the current study.

| AED       | Sample          | Location            | Reference     |
|-----------|-----------------|---------------------|---------------|
| 0.81      | Concrete blocks | Jos, Nigeria        | [17]          |
| 1.20      | Soil            | Jos, Nigeria        | [18]          |
| 0.33      | Sand            | Pakistan            | [33]          |
| 0.68      | Soil            | India               | [34]          |
| 0.40-0.83 | Sediments       | South-west, Nigeria | [35]          |
| 0.53      | Soil            | Iran                | [36]          |
| 0.88      | Sediment        | China               | [37]          |
| 0.36      | Tailings        | Nasarawa, Nigeria   | [38]          |
| 0.11      | Sediment        | Italy               | [39]          |
| 0.52      | Tailings        | Jos, Nigeria        | Present study |
| p = 0.430 |                 |                     |               |

Table 8. estimated annual gonadal dose equivalent, external and internal hazard indices, and excess life cancer risks from studied samples.

| Sample         | AGDE ( $\mu$ Sv/y) | $H_{ex}$ | $H_{in}$ | $ELCR_{out} \times 10^{-3}$ | $ELCR_{in} \times 10^{-3}$ |
|----------------|--------------------|----------|----------|-----------------------------|----------------------------|
| KK1            | 392.26             | 0.33     | 0.60     | 0.248                       | 1.953                      |
| KK2            | 595.18             | 0.51     | 0.94     | 0.377                       | 2.981                      |
| KK3            | 365.60             | 0.31     | 0.56     | 0.231                       | 1.818                      |
| KK4            | 365.57             | 0.31     | 0.52     | 0.229                       | 1.793                      |
| KK5            | 157.79             | 0.13     | 0.19     | 0.097                       | 0.750                      |
| KJ1            | 282.76             | 0.23     | 0.38     | 0.175                       | 1.375                      |
| KJ2            | 426.86             | 0.37     | 0.62     | 0.267                       | 2.101                      |
| KJ3            | 275.27             | 0.22     | 0.34     | 0.169                       | 1.317                      |
| KJ4            | 427.10             | 0.35     | 0.57     | 0.264                       | 2.068                      |
| KJ5            | 467.85             | 0.39     | 0.68     | 0.293                       | 2.302                      |
| KJ6            | 1479.90            | 1.29     | 2.28     | 0.938                       | 7.356                      |
| ZS1            | 205.37             | 0.18     | 0.33     | 0.131                       | 1.029                      |
| ZS2            | 169.42             | 0.14     | 0.22     | 0.105                       | 0.820                      |
| ZS3            | 150.76             | 0.13     | 0.21     | 0.094                       | 0.733                      |
| ZS4            | 177.61             | 0.16     | 0.28     | 0.113                       | 0.887                      |
| ZS5            | 378.93             | 0.32     | 0.57     | 0.238                       | 1.875                      |
| ZS6            | 367.35             | 0.31     | 0.52     | 0.230                       | 1.801                      |
| ZS7            | 281.10             | 0.24     | 0.39     | 0.176                       | 1.372                      |
| ZS8            | 228.53             | 0.19     | 0.34     | 0.144                       | 1.129                      |
| ZS9            | 251.73             | 0.21     | 0.35     | 0.157                       | 1.231                      |
| ZS10           | 318.55             | 0.27     | 0.44     | 0.199                       | 1.554                      |
| Mean           | 369.78             | 0.31     | 0.54     | 0.232                       | 1.821                      |
| Refrence value | 300.00             | 1.00     | 1.00     | 0.290                       | 0.290                      |
| P              | 0.265              | 0.001    | 0.001    | 0.150                       | 0.001                      |

The estimated annual gonadal dose equivalent AGDE from this study ranges from 150.76 to 1479.90  $\mu$ Sv/y with a mean value of 369.78  $\mu$ Sv/y (table 8).  $H_{ex}$  ranges from 0.13 to 1.29 with a mean of 0.31, and  $H_{in}$  ranges from 0.19 to 2.28 with a mean of 0.54. Obtained values for outdoor and indoor excess life cancer risks ( $ELCR_{out}$  and  $ELCR_{in}$ ) range from 0.094 to  $0.938 \times 10^{-3}$  and 0.733 to  $7.356 \times 10^{-3}$  with mean values of 0.232 and  $1.821 \times 10^{-3}$ , respectively.

### DISCUSSION

Tables 2 to 4 present a comparative analysis of the specific activities of radionuclides <sup>238</sup>U, <sup>232</sup>Th and <sup>40</sup>K from samples collected in the study areas against global averages. The findings reveal significant

variations, particularly in the concentrations of  $^{238}\text{U}$ . Samples from Kuru-Jenta show notably high mean specific activity for  $^{238}\text{U}$ , averaging 125.30 Bq/kg, which is substantially above the global average of 33 Bq/kg. In tailings from Kanikon, the mean specific activity of  $^{238}\text{U}$  is about three times higher than the global average. Sample KJ6 from Kuru-Jenta exhibits an exceptional increase, with  $^{238}\text{U}$  levels roughly 11 times above the global average. The overall mean concentration in Kuru-Jenta is over four times higher than the global standard.

Although the mean specific activity of tailings from Zagong-soil (table 4) is lower than the values for tailings for Kuru-Jenta, about 80% of the samples from Zangong-soil still show  $^{238}\text{U}$  concentrations that exceed the world's average. The mean specific activity obtained for  $^{238}\text{U}$  in these samples is about 70% higher than the world's average, while the mean values for  $^{232}\text{Th}$  and  $^{40}\text{K}$  are 79% and 69% lower than their respective world's average values in soil. Generally, the specific activities of  $^{232}\text{Th}$  and  $^{40}\text{K}$  across all samples are lower than global averages of 45 and 420 Bq/kg, respectively (28). For instance, tailings from Kanikon show mean values approximately 82% lower for  $^{232}\text{Th}$  and 65% lower for  $^{40}\text{K}$ . In Kuru-Jenta, the specific activities of  $^{232}\text{Th}$ , and  $^{40}\text{K}$  are about 59% and 32% lower than the global averages. Tailings from Zagong-Soil also show lower concentrations for  $^{232}\text{Th}$  and  $^{40}\text{K}$  compared to Kuru-Jenta.

Statistical analysis, represented by the p-values, indicates that the deviations from world averages are significant for  $^{238}\text{U}$  in some locations ( $p < 0.05$ ), while  $^{232}\text{Th}$  and  $^{40}\text{K}$  generally show lower levels of significance. The specific activities of  $^{238}\text{U}$ ,  $^{232}\text{Th}$ , and  $^{40}\text{K}$  observed in this study are comparable to those reported for other mining fields (table 5). However, some variations in data have been noted, with the specific activity values in this study being lower than those reported in earlier studies from the same region (8, 30, 31). Despite this, the mean  $^{238}\text{U}$  value from this study is higher than those reported by (25, 32), although those studies recorded higher values for  $^{40}\text{K}$ . The variations may be due to differences in mining intensity, the types of mined minerals, as well as climatic conditions and erosion levels during sample collection. Additionally, the solubility of certain radionuclides in water could contribute to the differences in observed values.

The radiological hazard indices estimated in Table 6 show that the  $R_{\text{eq}}$  and  $D_{\text{out}}$ , and  $AED_{\text{out}}$  are lower than their corresponding world's average and reference values by about 32%, 5.21%, and 17.11%, respectively. However, certain samples still exceed safety thresholds, raising concerns about their use as building materials. The mean indoor absorbed dose  $D_{\text{in}}$  is about 26.19% higher than the population-weighted average value of 84 nGy/h (28), with the highest value obtained being about four times higher

than the world's average value. Similarly, the mean annual effective dose indoors ( $AED_{\text{in}}$ ) is approximately 24% higher than the global average of 420  $\mu\text{Sv}/\text{y}$ , and 52.4% of the samples have values exceeding this average, with the highest being five times the world average. These results further reveal that tailings are not materials suitable for building.

The  $AED_{\text{in}}$  values obtained from this study are comparable to those reported for other building materials worldwide (table 7). The value from the present study is within the range of the values obtained by some similar studies (35, 36), above the results obtained by others (34, 38, 39), but lower than the results of some others (17, 18, 34, 37).

The radiological hazard indices indicate that while the mean values for the internal and external hazard indices ( $H_{\text{ex}}$  and  $H_{\text{in}}$ ) remain within acceptable limits, certain samples exceed these thresholds. The effective dose equivalents ( $AED_{\text{in}}$  and  $AGDE$ ) are higher than global averages, implying potential genetic radiation effects and an increased risk of cancer for individuals exposed to these tailings. The estimated excess lifetime cancer risk (ELCR) for outdoor exposure is about 80% of the total ELCR for all carcinogens, while the indoor ELCR is approximately six times higher than the reference value for all carcinogens. The ELCR values for residents living in homes built with these tailings are particularly concerning, indicating a significantly higher probability of cancer compared to global averages.

The mean values of  $ELCR_{\text{out}}$  and  $ELCR_{\text{in}}$  from this study are  $0.232 \times 10^{-3}$  and  $1.82110^{-3}$ , respectively. This suggests that mining activities in Kuru District add an additional risk of 232 cancer cases per million people for outdoor exposure and 1821 cancer cases per million for indoor exposure in homes constructed with these tailings. The elevated levels of  $^{238}\text{U}$ , particularly in Kuru-Jenta, present significant health hazards. Prolonged exposure to uranium is known to cause damage to various organs, including the cardiovascular system, liver, kidneys, and nervous system, and increase the risk of cancer (40, 41). This concern is heightened when uranium-contaminated tailings are used in construction. From a radiation protection standpoint, the primary concern is the increased likelihood of cancer over an individual's lifetime. The elevated specific activity of  $^{238}\text{U}$  in tailings from Kuru mining sites and the associated health risks underscore the unsuitability of these materials for construction and the potential significant rise in cancer risks to miners because of mining activities.

## CONCLUSION

The findings reveal that the mean specific activity of  $^{238}\text{U}$  in tailings, as well as mean values of  $D_{\text{in}}$ ,  $AED_{\text{in}}$ ,

and AGDE, and  $R_{aeq}$ ,  $D_{out}$ , and  $AED_{out}$  for some samples exceed global average or reference values. The elevated specific activity of  $^{238}\text{U}$  indicates an increased risk of internal radiation exposure for miners and residents primarily due to inhalation. The study also estimates an additional cancer risk of 2053 cases per 1 million people as derived from the Excess Lifetime Cancer Risk (ELCR) directly linked to mining activities in the Kuru District. From a radiation protection perspective, both the mining operations and the use of tailings as construction materials pose significant radiological risks. The heightened cancer risk underscores the need for enhanced safety measures and alternative materials to reduce these risks.

#### ACKNOWLEDGMENT

None.

**Ethical consideration:** None.

**Funding:** No funding was provided for this work.

**Data Availability:** Not Applicable.

**Conflict of Interest:** The authors declare that they have no competing interest in this study.

**Author Contributions:** A.M.B., contributed to the concept and design of the study, data collection analysis and interpretation, and wrote part of the manuscript. I.A.Z., contributed to the design of the study, wrote part of the manuscript, and reviewed and edited the first draft of the manuscript critically for intellectual content. O.K.I., took part in data analysis and interpretation. All authors read and approved the final manuscript.

#### REFERENCES

- International Atomic Energy Agency (IAEA), (2004) Radiation, people, and the environment (INIS-XA-703). Ford, J. (Ed.). IAEA, Vienna.
- Jwanbot DI, Izam MM, Nyam GG (2012) Radioactivity in some food crops from a high background radiation area on the Jos-Plateau, Nigeria. *J Nat Sci Res*, **2**(6): 76-79.
- Kolkoma D, Pereira F, Jojo PJ (2020) Assessment of radiological exposures in the vicinity of the gold mining area of Wau-Bulolo in Papua New Guinea. *Int J Radiat Res*, **20**(4): 773-777.
- Abd-Elkader MM, Shinonaga T, Sherif MM (2021). Radiological hazard assessments of radionuclides in building materials, soils, and sands from the Gaza Strip and the north of the Sinai Peninsula. *Sci Rep*, **11**: 23251.
- Nhan DD, Fernando CP, Ha NT, et al. (2012) Radon ( $^{222}\text{Rn}$ ) concentration in indoor air near the coal mining area of Nui Beo, North of Vietnam. *J Environ Radioact*, **110**: 98-103.
- Zeeb H, and Shannoun F, (2009) WHO handbook on indoor radon: a public health perspective. World Health Organisation, Geneva.
- Arogunjo AM, Ohenhen HO, Olowookere SP (2004) A re-evaluation of occupancy factors for effective dose estimate in tropical environments. *Radiat Prot Dosimetry*, **112**(2): 259-265.
- Ademola JA (2008) Exposure to high background radiation level in the tin mining area of Jos Plateau, Nigeria. *J Radiat Prot*, **28**: 93-99.
- Mallo SJ (2012) The socio-economic impact of artisanal mining in Kuru (Naraguta sheet 168) Plateau State, North Central Nigeria. *C J Eng Sci*, **7**(3): 27-32.
- Canavesio R (2014) Formal mining investments and artisanal mining in southern Madagascar: Effects of spontaneous reactions and adjustment policies on poverty alleviation. *Land Use Policy*, **36**: 145-154.
- Jibiri NN, Farai IP, Alausa SK (2007) Activity concentration of  $^{226}\text{Ra}$ ,  $^{232}\text{Th}$ , and  $^{40}\text{K}$  in different food crops from a high background radiation area in Bisichi, Jos Plateau, Nigeria. *Radiat Environ Biophys*, **46**(1): 53-59.
- Aliyu AS, Mousseau TA, Garba NN, et al. (2015) Estimation of annual effective dose due to ingestion of natural radionuclides in cattle in tin mining areas of Jos Plateau, Nigeria: Are large mammals really affected? *Nat Sci*, **7**: 190-196.
- Melodi M and Mary (2020) Productivity of artisanal tin mining in Jos South Local Government Area, Plateau State, Nigeria. *Int J Engin Adv Technol*, **8**(1): 46-54.
- Ugwu LU, Bashiru L, Okoli, JC (2020) Measurement of environmental radioactivity in mining dumpsites in Jos Metropolis, Plateau State, Nigeria. *J Appl Phys*, **12**(4): 25-28.
- Oresegun MO and Babalola IA (1988) Annual indoor dose burden estimates in dwellings built in Nigeria with radioactive U-Th rich tailings. Proceedings of the International Conference of Radiation Protection in Nuclear Energy International Atomic Energy Agency Sydney, Australia. IAEA-CN-51/7, **11**: 159-166.
- Atipo M, Olarinoye O, Awojogbe B, et al. (2020) High Terrestrial Radiation Level in an Active Tin-Mine at Jos South, Nigeria. *J Appl Sci Environ Manag*, **24** (3): 435-442.
- Ademola JA and Farai IP (2006) Gamma activity and radiation dose in concrete building blocks used for construction of dwellings in Jos, Nigeria. *Radiat Protect Dosimetry*, **121**: 395-398.
- Abba HT, Saleh MA, Hassan WMSW, et al. (2017) Mapping of natural gamma radiation (NGR) dose rate distribution in tin mining areas of Jos Plateau, Nigeria. *Enviro Earth Sci*, **76**(208): 1-9.
- Alausa SK (2020) Radiometric assessment of farm soils and food crops grown in Kuru-Jos, Nigeria. *Iran J Med Phys*, **17**: 289-297.
- Adedokun MB, Aweda MA, Maleka PP, et al. (2019) Natural radioactivity contents in commonly consumed leafy vegetables cultivated through surface water irrigation in Lagos State. *J Radiat Res Appl Sc*, **12**(1): 147-156.
- Currie LA (1968) Limits for qualitative detection and quantitative determination. *Anal Chem*, **40**: 586-593.
- Adedokun MB, Aweda MA, Maleka P, et al. (2022) Evaluation of natural radionuclides and associated radiation hazard indices in soil and water from selected vegetable farmlands in Lagos, Nigeria. *Environ Forensics*, **23**(3-40): 301-313.
- Jahan I, Ali M, Haydar MA, et al. (2016) Distribution of natural and probable artificial radioactivity in the sediment and water samples collected from low-lying areas of the Savar industrial zone, Bangladesh. *J Nucl Part Phys*, **6**(2): 25-34.
- Agalga R, Darko EO, Schandorf C (2013) Preliminary study on the levels of natural radionuclides in sediments of the Tono irrigation dam, Navrongo. *Int J Sci Technol*, **2**(11): 770-773.
- Ademola AK, Bello AK, Adeniyi AC (2014) Determination of natural radioactivity and hazards in soil samples in and around a gold mining area in Itangunmodi, South-Western Nigeria. *J Radiat Res Appl Sci*, **7**(3): 249-255.
- Qureshi AA, Tariq S, Din KU, et al. (2014). Evaluation of excessive lifetime cancer risk due to natural radioactivity in the river sediments of Northern Pakistan. *J Radiat Res Appl Sci*, **7**(4): 438-447.
- Orosun MM, Usikalu MR, Oyewumi KJ (2020) Radiological hazards assessment of the laterite mining field in Ilorin, north-central Nigeria. *Int J Radiat Res*, **18**(4): 895-906.
- UNSCEAR, (2000). Annex B: Exposure from Natural Radiation Sources. Report to the General Assembly, *United Nations Scientific Committee on the Effects of Atomic Radiation* (UNSCEAR), New York.
- Kurnaz A, Kucukomeroglu B, Keser R, et al. (2007) Determination of radioactivity levels and hazards of soil and sediment samples in Firtina Valley (Rize, Turkey). *App Radiat Isot*, **65**(11): 1281-1289.
- Ibeanu I (1999) Assessment of radiological hazards of tin mining and ore processing in Jos, Nigeria (IAEA-SM-359). International Atomic Energy Agency (IAEA).
- Masok FB, Masiteng PL, Daniel JI (2015) Natural radioactivity concentration and effective dose rate from Jos Tin mining dumpsites in Rayfield, Nigeria. *J Environ Earth Sci*, **5**: 51-55.
- Ojo JO (2016) Radioactivity in some foods and soils from Jos Tin mines, Jos Nigeria *Ife J Sci*, **18**(4): 1065-1071.
- Rahman S, Matiullah S, Mujahid A, et al. (2008) Assessment of radiological hazards due to the presence of natural radionuclides in samples of building materials collected from the north-western areas of Pakistan. *J Radiol Prot*, **28**(2): 205-212.
- Ramola RC, Gusain GS, Badoni M, et al. (2008)  $^{226}\text{Ra}$ ,  $^{232}\text{Th}$ , and  $^{40}\text{K}$  contents in soil samples from Garhwal Himalaya, India, and its radiological implications. *J Radiol Prot*, **28**: 379-385.

35. Farai IP and Isinkaye MO (2009) Radiological safety assessment of surface-water dam sediments used as building material in southwestern Nigeria. *J Radiol Prot*, **29**(1): 85-93.
36. Pashazadeh AM, Aghajani M, Nabipour I, *et al.* (2014) Annual effective dose from environmental gamma radiation in Bushehr city. *J Environ Health Sci Eng*, **12**(1): 1-4.
37. Lu X, Pan H, Ren C, *et al.* (2016) Natural radioactivity in reservoir sediment near an industrial park of northwest China. *J Radiol Prot*, **36**(2): 26-33.
38. Rilwan U, Umar I, Onochukwu GC, *et al.* (2020) Evaluation of radiation hazard indices in mining sites of Nasarawa State, Nigeria. *Asian J Res Rev Phys*, **3**(1): 8-16.
39. Caridi F, DiBella M, Sabatino G, *et al.* (2021) Assessment of natural radioactivity and radiological risks in river sediments from Calabria (Southern Italy). *Appl Sci*, **11**(4): 1729.
40. Vrhovnik P, Serafimovski T, Smuc NR *et al.* (2014) Uranium in different samples from Eastern Macedonia-a case study. *Procedia Earth Planet Sci*, **8**: 98-102.
41. Keith S, Faroon O, Roney N, *et al.* (2013). Toxicological profile for uranium. Agency for toxic substances and disease registry (US).